arXiv:1210.3749v1 [quant-ph] 14 Oct 2012

Quantum Local Asymptotic Normality based on a new Quantum Likelihood Ratio

Koichi Yamagata, Akio Fujiwara
Department of Mathematics, Osaka University
Toyonaka 560-0043, Osaka, Japan
and
Richard D. Gill
Mathematical Institute, Faculty of Science
Leiden University

October 16, 2012

Abstract

We develop a theory of local asymptotic normality in a quantum regime based on a novel quantum analogue of the log-likelihood ratio. This formulation is applicable to any quantum statistical model satisfying a mild smoothness condition. As an application, we prove the asymptotic achievability of the Holevo bound for the local shift parameter.

1 Introduction

Given a (classical) statistical model $\mathcal{S} = \{p_\theta\, ;\, \theta \in \Theta\}$ on a probability space $(\Omega, \mathcal{F}, \mu)$ indexed by a parameter θ that ranges over an open subset Θ of $\mathbb{R}^d$, let us introduce a local parameter $h := \sqrt{n}(\theta - \theta_0)$ around a fixed $\theta_0 \in \Theta$. If the parametrization $\theta \mapsto p_\theta$ is sufficiently smooth, it is known that the statistical properties of the model $\left\{p_{\theta_0+h/\sqrt{n}}^{\otimes n}\, ;\, h \in \mathbb{R}^d\right\}$ is similar to that of the Gaussian shift model $\left\{N(h, J_{\theta_0}^{-1})\, ;\, h \in \mathbb{R}^d\right\}$ for large n, where $p_\theta^{\otimes n}$ is the nth i.i.d. extension of p_θ, and J_{θ_0} is the Fisher information matrix of the model p_θ at θ_0. This property is called the local asymptotic normality of the model $\mathcal{S}$ [20].

More generally, a sequence $\left\{p_\theta^{(n)}\, ;\, \theta \in \Theta \subset \mathbb{R}^d\right\}$ of statistical models on $(\Omega^{(n)}, \mathcal{F}^{(n)}, \mu^{(n)})$ is called *locally asymptotically normal* (LAN) at $\theta_0 \in \Theta$ if there exist a $d \times d$ positive matrix J and random vectors $\Delta^{(n)} = (\Delta_1^{(n)}, \ldots, \Delta_d^{(n)})$ such that $\Delta^{(n)} \overset{0}{\rightsquigarrow} N(0, J)$ and

$$\log \frac{p_{\theta_0+h/\sqrt{n}}^{(n)}}{p_{\theta_0}^{(n)}} = h^i \Delta_i^{(n)} - \frac{1}{2} h^i h^j J_{ij} + o_{p_{\theta_0}}(1)$$

for all $h \in \mathbb{R}^d$. Here the arrow $\overset{h}{\rightsquigarrow}$ stands for the convergence in distribution under $p_{\theta_0+h/\sqrt{n}}^{(n)}$, the remainder term $o_{p_{\theta_0}}(1)$ converges in probability to zero under $p_{\theta_0}^{(n)}$, and Einstein's summation convention is used. The above expansion is similar in form to the log-likelihood ratio of the Gaussian shift model:

$$\log \frac{dN(h, J^{-1})}{dN(0, J^{-1})}(X^1, \ldots, X^d) = h^i (X^j J_{ij}) - \frac{1}{2} h^i h^j J_{ij}.$$

This is the underlying mechanism behind the statistical similarities between models $\left\{p_{\theta_0+h/\sqrt{n}}^{(n)}\,;\ h\in\mathbb{R}^d\right\}$ and $\left\{N(h,J^{-1})\,;\ h\in\mathbb{R}^d\right\}$.

In order to put the similarities to practical use, one needs some mathematical devices. In general, a statistical theory comprises two parts. One is to prove the existence of a statistic that possesses a certain desired property (direct part), and the other is to prove the non-existence of a statistic that exceeds that property (converse part). In the problem of asymptotic efficiency, for example, the converse part, the impossibility to do asymptotically better than the best which can be done in the limit situation, is ensured by the following proposition, which is usually referred to as "Le Cam's third lemma" [20].

Proposition 1. *Suppose* $\left\{p_\theta^{(n)}\,;\ \theta\in\Theta\subset\mathbb{R}^d\right\}$ *is LAN at* $\theta_0\in\Theta$*, with* $\Delta^{(n)}$ *and* J *being as above, and let* $X^{(n)}=(X_1^{(n)},\dots,X_r^{(n)})$ *be a sequence of random vectors. If the joint distribution of* $X^{(n)}$ *and* $\Delta^{(n)}$ *converges to a Gaussian distribution, in that*

$$\begin{pmatrix}X^{(n)}\\ \Delta^{(n)}\end{pmatrix}\overset{0}{\rightsquigarrow}N\left(\begin{pmatrix}0\\0\end{pmatrix},\begin{pmatrix}\Sigma & \tau\\ {}^t\tau & J\end{pmatrix}\right),$$

then $X^{(n)}\overset{h}{\rightsquigarrow}N(\tau h,\Sigma)$ *for all* $h\in\mathbb{R}^d$.

Now, it appears from this lemma that it already tells us something about the direct problem. In fact, by putting $X^{(n)j}:=\sum_{k=1}^d\left[J^{-1}\right]^{jk}\Delta_k^{(n)}$, we have

$$\begin{pmatrix}X^{(n)}\\ \Delta^{(n)}\end{pmatrix}\overset{0}{\rightsquigarrow}N\left(\begin{pmatrix}0\\0\end{pmatrix},\begin{pmatrix}J^{-1} & I\\ I & J\end{pmatrix}\right),$$

so that $X^{(n)}\overset{h}{\rightsquigarrow}N(h,J^{-1})$ follows from Proposition 1. This proves the existence of an asymptotically efficient estimator for h. In the real world however, we do not know θ_0 (obviously!). Thus the existence of an asymptotically optimal estimator for h does not translate into the existence of an asymptotically optimal estimator of θ. In fact, the usual way that Le Cam's third lemma is used in the subsequent analysis is in order to prove the so-called representation theorem, [20, Theorem 7.10]. This theorem can be used to tell us in several precise mathematical senses that no estimator can asymptotically do better than what can be achieved in the limiting Gaussian model.

For instance, Van der Vaart's version of the representation theorem leads to the asymptotic minimax theorem, telling us that the worst behaviour of an estimator as θ varies in a shrinking (1 over root n) neighbourhood of θ_0 cannot improve on what we expect from the limiting problem. This theorem applies to *all* possible estimators, but only discusses their *worst* behaviour in a neighbourhood of θ. Another option is to use the representation theorem to derive the convolution theorem, which tells us that *regular* estimators (estimators whose asymptotic behaviour in a small neighbourhood of θ is more or less stable as the parameter varies) have a limiting distribution which in a very strong sense is more disperse than the optimal limiting distribution which we expect from the limiting statistical problem.

This paper addresses a quantum extension of LAN (abbreviated as QLAN). As in the classical statistics, one of the important subjects of QLAN is to show the existence of an estimator (direct part) that enjoys certain desired properties. Some earlier works of asymptotic quantum parameter estimation theory revealed the asymptotic achievability of the Holevo bound, a quantum extension of the Cramér-Rao type bound (cf., Appendices A, B). Using a group representation theoretical method, Hayashi and Matsumoto [8] showed that the Holevo bound for the quantum statistical model $\mathcal{S}(\mathbb{C}^2)=\{\rho_\theta\,;\,\theta\in\Theta\}$ comprising the totality of density operators on the Hilbert space $\mathcal{H}\simeq\mathbb{C}^2$ is asymptotically achievable at a given single point $\theta_0 \in \Theta$. Following their work, Guţă and Kahn [5, 11] developed a theory of strong QLAN, and proved that the Holevo bound is asymptotically uniformly achievable around a given $\theta_0 \in \Theta$ for the quantum statistical model $\mathcal{S}(\mathbb{C}^D) = \{\rho_\theta\,;\,\theta \in \Theta\}$ comprising the totality of density operators on the finite dimensional Hilbert space $\mathcal{H} \simeq \mathbb{C}^D$. They proved that $\left\{\rho_{h/\sqrt{n}}^{\otimes n}\,;\, h \in \mathbb{R}^d\right\}$ and a quantum Gaussian shift model can be translated by quantum channels to each other asymptotically. Although their result is powerful, their QLAN has several drawbacks. First of all, their method works only for i.i.d. extension of the totality $\mathcal{S}(\mathcal{H})$ of the quantum states on the Hilbert space $\mathcal{H}$, and is not applicable to generic submodels of $\mathcal{S}(\mathcal{H})$. Moreover, it makes use of a special parametrization θ of $\mathcal{S}(\mathcal{H})$, in which the change of eigenvalues and eigenvectors are treated as essential. Furthermore, it does not work if the reference state ρ_{θ_0} has a multiplicity of eigenvalues. Hayashi and Matsumoto's formulation [8] also suffers from the same problems.

The purpose of the present paper is to develop a theory of (weak) QLAN based on a new quantum extension of the log-likelihood ratio. This formulation is applicable to any quantum statistical model satisfying a mild smoothness condition, and is free from artificial setups such as the use of a special coordinate system and/or non-degeneracy of eigenvalues of the reference state at which QLAN works. We also prove asymptotic achievability of the Holevo bound for the local shift parameter h that belong to a dense subset of $\mathbb{R}^d$.

This paper is organized as follows. The main results are summarized in Section 2. We first introduce a novel type of quantum log-likelihood ratio, and define a quantum extension of local asymptotic normality in a quite analogous way to the classical LAN. We then explore some basic properties of QLAN, including a sufficient condition for an i.i.d. model to be QLAN, and a quantum extension of Le Cam's third lemma. Proofs of those results are provided in Section 3. Section 4 is devoted to application of QLAN, including the asymptotic achievability of the Holevo bound and asymptotic estimation theory for some typical qubit models. For the reader's convenience, a brief account of quantum estimation theory are presented in appendices A-D. Those prerequisites are used throughout the paper.

It is also important to notice the limits of this work, which means that there are many open problems left to study in the future. In the classical case, the theory of LAN builds, of course, on the rich theory of convergence in distribution, as studied in probability theory. In the quantum case, there still does not exist a full parallel theory. Some of the most useful lemmas in the classical theory simply are not true when translated in the quantum domain. For instance, in the classical case, we know that if the sequence of random variables X_n converges in distribution to a random variable X, and at the same time the sequence Y_n converges in probability to a constant c, then this implies joint convergence in distribution of (X_n, Y_n) to the pair (X, c). The obvious analogue of this in the quantum domain is simply untrue. In fact, there is not even a general theory of convergence in distribution at all: there is only a theory of convergence in distribution towards quantum Gaussian limits. Unfortunately, even in this special case the natural analogue of the just mentioned result simply fails to be true.

Because of these obstructions we are not at present able to follow the standard route from Le Cam's third lemma to the representation theorem, and from there to asymptotic minimax or convolution theorems.

However we believe that the paper presents some notable steps in this direction. Moreover, just as with Le Cam's third lemma, one is able to use the lemma to construct what can be conjectured to be asymptotically optimal measurement and estimation schemes. We make some more remarks on these possibilities later in the paper.

2 Main results

Definition 2 (Quantum log-likelihood ratio)**.** We say a pair of density operators ρ and σ on a finite dimensional Hilbert space $\mathcal{H}$ are *mutually absolutely continuous,* $\rho \sim \sigma$ in symbols, if there exist a Hermitian operator $\mathcal{L}$ that satisfies

$$\sigma = \mathrm{e}^{\frac{1}{2}\mathcal{L}} \rho\, \mathrm{e}^{\frac{1}{2}\mathcal{L}}.$$

We shall call such a Hermitian operator $\mathcal{L}$ a *quantum log-likelihood ratio.* When the reference states ρ and σ need to be specified, $\mathcal{L}$ shall be denoted by $\mathcal{L}\left(\sigma|\rho\right)$, so that

$$\sigma = \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)} \rho\, \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}.$$

We use the convention that $\mathcal{L}\left(\rho|\rho\right) = 0$.

Example 3. Faithful states are always mutually absolutely continuous. In fact, given $\rho > 0$ and $\sigma > 0$, they are related as $\sigma = \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}\rho \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}$, where

$$\mathcal{L}\left(\sigma|\rho\right) = 2\log\left(\sqrt{\rho^{-1}}\sqrt{\sqrt{\rho}\sigma\sqrt{\rho}}\sqrt{\rho^{-1}}\right) = 2\log\left(\sqrt{\sigma}\left(\sqrt{\sqrt{\sigma}\rho\sqrt{\sigma}}\right)^{-1}\sqrt{\sigma}\right).$$

Note that $\mathrm{Tr}\,\rho\, \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}$ is identical to the fidelity between ρ and σ, and $\mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}$ is nothing but the operator geometric mean $\sigma\#\rho^{-1}$, where $A\#B := A^{1/2}\left(A^{-1/2}BA^{-1/2}\right)^{1/2}A^{1/2}$ for positive operators A, B.

Example 4. Pure states $\rho = \left|\psi\right\rangle\left\langle\psi\right|$ and $\sigma = \left|\xi\right\rangle\left\langle\xi\right|$ are mutually absolutely continuous if and only if $\left\langle\xi|\psi\right\rangle \neq 0$.

Proof. Suppose first that $\rho \sim \sigma$. Then

$$\left|\left\langle\xi|\psi\right\rangle\right|^2 = \mathrm{Tr}\,\rho\sigma = \mathrm{Tr}\,\left|\psi\right\rangle\left\langle\psi\right| \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}\left|\psi\right\rangle\left\langle\psi\right| \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)} = \left|\left\langle\psi\right| \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}\left|\psi\right\rangle\right|^2 > 0.$$

Suppose next that $\left\langle\xi|\psi\right\rangle \neq 0$. Then

$$R := I + \frac{1}{\left|\left\langle\xi|\psi\right\rangle\right|}\left|\xi\right\rangle\left\langle\xi\right| - \left|\psi\right\rangle\left\langle\psi\right|$$

is positive definite, and $\mathcal{L}\left(\sigma|\rho\right) := 2\log R$ satisfies

$$\mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}\left|\psi\right\rangle = R\left|\psi\right\rangle = \frac{\left\langle\xi|\psi\right\rangle}{\left|\left\langle\xi|\psi\right\rangle\right|}\left|\xi\right\rangle,$$

showing that $\rho \sim \sigma$. Note that $\mathrm{Tr}\,\rho\, \mathrm{e}^{\frac{1}{2}\mathcal{L}(\sigma|\rho)}$ is the fidelity again. □

Given a $d \times d$ real skew-symmetric matrix S, let $\mathrm{CCR}\left(S\right)$ be the CCR algebra defined by

$$\mathrm{e}^{\sqrt{-1}X_i}\mathrm{e}^{\sqrt{-1}X_j} = \mathrm{e}^{\sqrt{-1}S_{ij}}\mathrm{e}^{\sqrt{-1}(X_i+X_j)} \quad (1 \leq i, j \leq d),$$

(see [14, 18, 10, 9]). We call $X = (X_1, \ldots, X_d)$ the basic canonical observables of $\mathrm{CCR}\left(S\right)$. A state ϕ on $\mathrm{CCR}\left(S\right)$ is characterized by the *characteristic function* $\mathcal{F}_\xi\{\phi\} := \phi(\mathrm{e}^{\sqrt{-1}\xi^i X_i})$, where $\xi = (\xi^i)_{i=1}^d \in \mathbb{R}^d$ and Einstein's summation convention is used. A state ϕ on $\mathrm{CCR}\left(S\right)$ is called a *quantum Gaussian state*, denoted by $\phi \sim N(h, J)$, if the characteristic function takes the form

$$\mathcal{F}_\xi\{\phi\} = \mathrm{e}^{\sqrt{-1}\xi^i h_i - \frac{1}{2}\xi^i\xi^j V_{ij}},$$

where $h = (h_i)_{i=1}^d \in \mathbb{R}^d$ and $V = (V_{ij})$ is a real symmetric matrix such that the Hermitian matrix $J := V + \sqrt{-1}S$ is positive semidefinite. When the canonical observables X need to be specified, we also use the notation $(X, \phi) \sim N(h, J)$.

We will discuss relationships between a quantum Gaussian state ϕ on a CCR and a state on another algebra. In such a case, we need to use the *quasi-characteristic function*

$$\phi\left(\prod_{t=1}^{r} \mathrm{e}^{\sqrt{-1}\xi_t^i X_i}\right) = \exp\left(\sum_{t=1}^{r}\left(\sqrt{-1}\xi_t^i h_i - \frac{1}{2}\xi_t^i \xi_t^j J_{ji}\right) - \sum_{t=1}^{r}\sum_{s=t+1}^{r} \xi_t^i \xi_s^j J_{ji}\right), \tag{1}$$

of a quantum Gaussian state, where $(X, \phi) \sim N(h, J)$ and $\{\xi_t\}_{t=1}^r$ is a finite subset of $\mathbb{C}^d$ [10].

Given a sequence $\mathcal{H}^{(n)}$, $n \in \mathbb{N}$, of finite dimensional Hilbert spaces, let $X^{(n)} = (X_1^{(n)}, \dots, X_d^{(n)})$ and $\rho^{(n)}$ be a list of observables and a density operator on each $\mathcal{H}^{(n)}$. We say the sequence $\left(X^{(n)}, \rho^{(n)}\right)$ *converges in law to a quantum Gaussian state* $N(h, J)$, denoted as $(X^{(n)}, \rho^{(n)}) \underset{q}{\rightsquigarrow} N(h, J)$, if

$$\lim_{n\to\infty} \operatorname{Tr} \rho^{(n)} \left(\prod_{t=1}^{r} \mathrm{e}^{\sqrt{-1}\xi_t^i X_i^{(n)}}\right) = \phi\left(\prod_{t=1}^{r} \mathrm{e}^{\sqrt{-1}\xi_t^i X_i}\right)$$

for any finite subset $\{\xi_t\}_{t=1}^r$ of $\mathbb{C}^d$, where $(X, \phi) \sim N(h, J)$. Here we do not intend to introduce the notion of "quantum convergence in law" in general. We use this notion only for quantum Gaussian states in the sense of convergence of quasi-characteristic function.

The following is a version of the quantum central limit theorem (see [10], for example).

Proposition 5 (Quantum central limit theorem)**.** *Let A_i $(1 \le i \le d)$ and ρ be observables and a state on a finite dimensional Hilbert space $\mathcal{H}$ such that* $\operatorname{Tr} \rho A_i = 0$*, and let*

$$X_i^{(n)} := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes A_i \otimes I^{\otimes(n-k)}.$$

Then $(X^{(n)}, \rho^{\otimes n}) \underset{q}{\rightsquigarrow} N(0, J)$, where J is the Hermitian matrix whose (i,j)th entry is given by $J_{ij} = \operatorname{Tr} \rho A_j A_i$.

For later convenience, we introduce the notion of an "infinitesimal" object relative to the convergence $(X^{(n)}, \rho^{(n)}) \underset{q}{\rightsquigarrow} N(0, J)$ as follows. Given a list $X^{(n)} = (X_1^{(n)}, \dots, X_d^{(n)})$ of observables and a state $\rho^{(n)}$ on each $\mathcal{H}^{(n)}$ that satisfy $(X^{(n)}, \rho^{(n)}) \underset{q}{\rightsquigarrow} N(0, J) \sim (X, \phi)$, we say a sequence $R^{(n)}$ of observables, each being defined on $\mathcal{H}^{(n)}$, is *infinitesimal relative to the convergence* $(X^{(n)}, \rho^{(n)}) \underset{q}{\rightsquigarrow} N(0, J)$ if it satisfies

$$\lim_{n\to\infty} \operatorname{Tr} \rho^{(n)} \left(\prod_{t=1}^{r} \mathrm{e}^{\sqrt{-1}\left(\xi_t^i X_i^{(n)} + \eta_t R^{(n)}\right)}\right) = \phi\left(\prod_{t=1}^{r} \mathrm{e}^{\sqrt{-1}\xi_t^i X_i}\right) \tag{2}$$

for any finite subset of $\{\xi_t\}_{t=1}^r$ of $\mathbb{C}^d$ and any finite subset $\{\eta_t\}_{t=1}^r$ of $\mathbb{C}$. This is equivalent to saying that

$$\left(\begin{pmatrix} X^{(n)} \\ R^{(n)} \end{pmatrix}, \rho^{(n)}\right) \underset{q}{\rightsquigarrow} N\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} J & 0 \\ 0 & 0 \end{pmatrix}\right),$$

and is much stronger a requirement than

$$(R^{(n)}, \rho^{(n)}) \underset{q}{\rightsquigarrow} N(0, 0).$$

An infinitesimal object $R^{(n)}$ relative to $(X^{(n)}, \rho^{(n)}) \underset{q}{\rightsquigarrow} N(0, J)$ will be denoted as $o(X^{(n)}, \rho^{(n)})$.

The following is in essence a simple extension of Proposition 5, but will turn out to be useful in applications.

Lemma 6. *In addition to assumptions of Proposition 5, let $P(n)$, $n \in \mathbb{N}$, be a sequence of observables on $\mathcal{H}$, and let*

$$R^{(n)} := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes P(n) \otimes I^{\otimes(n-k)}.$$

If $\lim_{n\to\infty} P(n) = 0$ *and* $\lim_{n\to\infty} \sqrt{n}\, \mathrm{Tr}\, \rho P(n) = 0$, *then* $R^{(n)} = o(X^{(n)}, \rho^{\otimes n})$.

We are now ready to extend the notion of local asymptotic normality to a quantum regime.

Definition 7 (QLAN)**.** Given a sequence $\mathcal{H}^{(n)}$ of finite dimensional Hilbert spaces, let $\mathcal{S}^{(n)} = \left\{ \rho_\theta^{(n)} \,;\, \theta \in \Theta \subset \mathbb{R}^d \right\}$ be a quantum statistical model on $\mathcal{H}^{(n)}$, where $\rho_\theta^{(n)}$ is a parametric family of density operators and Θ is an open set. We say $\mathcal{S}^{(n)}$ is *quantum locally asymptotically normal* (QLAN) at $\theta_0 \in \Theta$ if the following conditions are satisfied:

(i) for any $\theta \in \Theta$ and $n \in \mathbb{N}$, $\rho_\theta^{(n)}$ is mutually absolutely continuous to $\rho_{\theta_0}^{(n)}$,

(ii) there exist a list $\Delta^{(n)} = (\Delta_1^{(n)}, \ldots, \Delta_d^{(n)})$ of observables on each $\mathcal{H}^{(n)}$ that satisfies

$$\left(\Delta^{(n)}, \rho_{\theta_0}^{(n)} \right) \underset{q}{\rightsquigarrow} N(0, J),$$

where J is a $d \times d$ Hermitian positive semidefinite matrix with $\mathrm{Re}\, J > 0$,

(iii) quantum log-likelihood ratio $\mathcal{L}_h^{(n)} := \mathcal{L}\left(\rho_{\theta_0 + h/\sqrt{n}}^{(n)} \middle| \rho_{\theta_0}^{(n)} \right)$ is expanded in $h \in \mathbb{R}^d$ as

$$\mathcal{L}_h^{(n)} = h^i \Delta_i^{(n)} - \frac{1}{2}(J_{ij} h^i h^j) I^{(n)} + o(\Delta^{(n)}, \rho_{\theta_0}^{(n)}), \tag{3}$$

where $I^{(n)}$ is the identity operator on $\mathcal{H}^{(n)}$.

It is also possible to extend Le Cam's third lemma (Proposition 1) to a quantum regime. To this end, however, we need a device to handle the infinitesimal residual term in (3) in a more elaborate way.

Definition 8. Let $\mathcal{S}^{(n)} = \left\{ \rho_\theta^{(n)} \,;\, \theta \in \Theta \subset \mathbb{R}^d \right\}$ be as in Definition 7, and let $X^{(n)} = (X_1^{(n)}, \ldots, X_r^{(n)})$ be a list of observables on $\mathcal{H}^{(n)}$. We say the pair $(\mathcal{S}^{(n)}, X^{(n)})$ is *jointly QLAN* at $\theta_0 \in \Theta$ if the following conditions are satisfied:

(i) for any $\theta \in \Theta$ and $n \in \mathbb{N}$, $\rho_\theta^{(n)}$ is mutually absolutely continuous to $\rho_{\theta_0}^{(n)}$,

(ii) there exist a list $\Delta^{(n)} = (\Delta_1^{(n)}, \ldots, \Delta_d^{(n)})$ of observables on each $\mathcal{H}^{(n)}$ that satisfies

$$\left(\begin{pmatrix} X^{(n)} \\ \Delta^{(n)} \end{pmatrix}, \rho_{\theta_0}^{(n)} \right) \underset{q}{\rightsquigarrow} N\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \Sigma & \tau \\ \tau^* & J \end{pmatrix} \right), \tag{4}$$

where Σ and J are Hermitian positive semidefinite matrices of size $r \times r$ and $d \times d$, respectively, with $\mathrm{Re}\, J > 0$, and τ is a complex matrix of size $r \times d$.

(iii) quantum log-likelihood ratio $\mathcal{L}_h^{(n)} := \mathcal{L}\left(\rho_{\theta_0 + h/\sqrt{n}}^{(n)} \middle| \rho_{\theta_0}^{(n)} \right)$ is expanded in $h \in \mathbb{R}^d$ as

$$\mathcal{L}_h^{(n)} = h^i \Delta_i^{(n)} - \frac{1}{2}(J_{ij} h^i h^j) I^{(n)} + o\left(\begin{pmatrix} X^{(n)} \\ \Delta^{(n)} \end{pmatrix}, \rho_{\theta_0}^{(n)} \right). \tag{5}$$

With Definition 8, we can state a quantum extension of Le Cam's third lemma as follows.

Theorem 9. *Let* $\mathcal{S}^{(n)}$ *and* $X^{(n)}$ *be as in Definition 8. If* $(\rho_\theta^{(n)}, X^{(n)})$ *is jointly QLAN at* $\theta_0 \in \Theta$*, then*

$$\left(X^{(n)},\, \rho_{\theta_0+h/\sqrt{n}}^{(n)}\right) \underset{q}{\rightsquigarrow} N((\operatorname{Re}\tau)\, h,\, \Sigma)$$

for any $h \in \mathbb{R}^d$*.*

In applications, we often handle i.i.d. extensions. In classical statistics, a sequence of i.i.d. extensions of a model is LAN if the log-likelihood ratio is twice differentiable [20]. Quite analogously, we can prove that a sequence of i.i.d. extensions of a quantum statistical model is QLAN if the quantum log-likelihood ratio is twice differentiable.

Theorem 10. *Let* $\left\{\rho_\theta \,;\, \theta \in \Theta \subset \mathbb{R}^d\right\}$ *be a quantum statistical model on a finite dimensional Hilbert space* $\mathcal{H}$ *satisfying* $\rho_\theta \sim \rho_{\theta_0}$ *for all* $\theta \in \Theta$*, where* $\theta_0 \in \Theta$ *is an arbitrarily fixed point. If* $\mathcal{L}_h := \mathcal{L}\left(\rho_{\theta_0+h} | \rho_{\theta_0}\right)$ *is differentiable around* $h = 0$ *and twice differentiable at* $h = 0$*, then* $\left\{\rho_\theta^{\otimes n} \,;\, \theta \in \Theta \subset \mathbb{R}^d\right\}$ *is QLAN at* θ_0*: that is,* $\rho_\theta^{\otimes n} \sim \rho_{\theta_0}^{\otimes n}$*, and*

$$\Delta_i^{(n)} := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes L_i \otimes I^{\otimes(n-k)}$$

and $J_{ij} := \operatorname{Tr} \rho_{\theta_0} L_j L_i$*, with* L_i *being the* i*th symmetric logarithmic derivative at* $\theta_0 \in \Theta$*, satisfy conditions (ii) (iii) in Definition 7.*

By combining Theorem 10 with Theorem 9, we obtain the following.

Corollary 11. *Let* $\left\{\rho_\theta \,;\, \theta \in \Theta \subset \mathbb{R}^d\right\}$ *be a quantum statistical model on* $\mathcal{H}$ *satisfying* $\rho_\theta \sim \rho_{\theta_0}$ *for all* $\theta \in \Theta$*, where* $\theta_0 \in \Theta$ *is an arbitrarily fixed point. Further, let* $\{B_i\}_{1 \le i \le r}$ *be observables on* $\mathcal{H}$ *satisfying* $\operatorname{Tr} \rho_{\theta_0} B_i = 0$ *for* $i = 1, \ldots, r$*. If* $\mathcal{L}_h := \mathcal{L}\left(\rho_{\theta_0+h} | \rho_{\theta_0}\right)$ *is differentiable around* $h = 0$ *and twice differentiable at* $h = 0$*, then the pair* $\left(\left\{\rho_\theta^{\otimes n}\right\},\, X^{(n)}\right)$ *of i.i.d. extension model* $\left\{\rho_\theta^{\otimes n}\right\}$ *and the list* $X^{(n)} = \{X_i^{(n)}\}_{1 \le i \le r}$ *of observables defined by*

$$X_i^{(n)} := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes B_i \otimes I^{\otimes(n-k)}$$

is jointly QLAN at θ_0*, and*

$$\left(X^{(n)}, \rho_{\theta_0+h/\sqrt{n}}^{\otimes n}\right) \underset{q}{\rightsquigarrow} N((\operatorname{Re}\tau)\, h, \Sigma)$$

for any $h \in \mathbb{R}^d$*, where* Σ *is the* $r \times r$ *positive semidefinite matrix defined by* $\Sigma_{ij} = \operatorname{Tr} \rho_{\theta_0} B_j B_i$ *and* τ *is the* $r \times d$ *matrix defined by* $\tau_{ij} = \operatorname{Tr} \rho_{\theta_0} L_j B_i$ *with* L_i *being the* i*th SLD at* θ_0*.*

As in the classical case, Corollary 11 prompts us to expect that any estimator for a quantum Gaussian shift model $\left\{N((\operatorname{Re}\tau)\, h, \Sigma)\,;\, h \in \mathbb{R}^d\right\}$ could be realized asymptotically on $\left\{\rho_{\theta_0+h/\sqrt{n}}^{\otimes n} \,;\, h \in \mathbb{R}^d\right\}$. This program will be partly demonstrated in Section 4 in the form of achievability of the Holevo bound.

3 Proofs of main theorems

3.1 Proof of Lemma 6

We shall prove (2) for $\{\xi_t\}_{t=1}^r \subset \mathbb{C}^d$ and $\{\eta_t\}_{t=1}^r \subset \mathbb{C}$.

$$\begin{aligned}
&\mathrm{Tr}\,\rho^{\otimes n}\left(\prod_{t=1}^r \mathrm{e}^{\sqrt{-1}\left(\xi_t^i X_i^{(n)}+\eta_t R^{(n)}\right)}\right)\\
&\quad= \mathrm{Tr}\,\rho^{\otimes n}\left[\prod_{t=1}^r \exp\left\{\frac{\sqrt{-1}}{\sqrt{n}}\sum_{k=1}^n I^{\otimes(k-1)}\otimes\left(\xi_t^i A_i+\eta_t P(n)\right)\otimes I^{\otimes(n-k)}\right\}\right]\\
&\quad= \mathrm{Tr}\,\rho^{\otimes n}\left[\prod_{t=1}^r \left\{\exp\left(\frac{\sqrt{-1}}{\sqrt{n}}\left(\xi_t^i A_i+\eta_t P(n)\right)\right)\right\}^{\otimes n}\right]\\
&\quad= \mathrm{Tr}\,\rho^{\otimes n}\left[\left\{\prod_{t=1}^r \exp\left(\frac{\sqrt{-1}}{\sqrt{n}}\left(\xi_t^i A_i+\eta_t P(n)\right)\right)\right\}^{\otimes n}\right]\\
&\quad= \left[\mathrm{Tr}\,\rho\left\{\prod_{t=1}^r \exp\left(\frac{\sqrt{-1}}{\sqrt{n}}\left(\xi_t^i A_i+\eta_t P(n)\right)\right)\right\}\right]^n\\
&\quad= \left[\mathrm{Tr}\,\rho\left\{\sum_{k_1,\dots,k_r\in\mathbb{Z}_+}\left(\frac{\sqrt{-1}}{\sqrt{n}}\right)^{k_1+\cdots+k_r}\prod_{t=1}^r\frac{1}{k_t!}\left(\xi_t^i A_i+\eta_t P(n)\right)^{k_t}\right\}\right]^n,
\end{aligned}$$

where $\mathbb{Z}_+ = \{0,1,2,\dots\}$. The terms corresponding to $k_1+\cdots+k_r=1$ in the summand are

$$\mathrm{Tr}\,\rho\left(\frac{\sqrt{-1}}{\sqrt{n}}\sum_{t=1}^r\left(\xi_t^i A_i+\eta_t P(n)\right)\right) = \left(\sum_{t=1}^r \eta_t\right)\frac{\sqrt{-1}}{\sqrt{n}}\mathrm{Tr}\,\rho P(n) = o\left(\frac{1}{n}\right)$$

because $\mathrm{Tr}\,\rho A_i = 0$ and $\mathrm{Tr}\,\rho P(n) = o(\frac{1}{\sqrt{n}})$. The terms corresponding to $k_1+\cdots+k_r=2$ are

$$\begin{aligned}
&-\frac{1}{n}\mathrm{Tr}\,\rho\left\{\sum_{k_1+\cdots+k_r=2}\left(\prod_{t=1}^r\frac{1}{k_t!}\left(\xi_t^i A_i+\eta_t P(n)\right)^{k_t}\right)\right\}\\
&\quad= -\frac{1}{2n}\sum_{t=1}^r \mathrm{Tr}\,\rho\left(\xi_t^i A_i+\eta_t P(n)\right)^2 - \frac{1}{n}\sum_{t=1}^r\sum_{s=t+1}^r \mathrm{Tr}\,\rho\left(\xi_t^i A_i+\eta_t P(n)\right)\left(\xi_s^j A_j+\eta_s P(n)\right)\\
&\quad= -\frac{1}{2n}\sum_{t=1}^r \xi_t^i\xi_t^j\,\mathrm{Tr}\,\rho A_i A_j - \frac{1}{n}\sum_{t=1}^r\sum_{s=t+1}^r \xi_t^i\xi_s^j\,\mathrm{Tr}\,\rho A_i A_j + o\left(\frac{1}{n}\right)\\
&\quad= -\frac{1}{2n}\sum_{t=1}^r \xi_t^i\xi_t^j J_{ji} - \frac{1}{n}\sum_{t=1}^r\sum_{s=t+1}^r \xi_t^i\xi_s^j J_{ji} + o\left(\frac{1}{n}\right).
\end{aligned}$$

In the third line, we used the fact that $P(n) = o(1)$. Let us denote the terms corresponding to $k_1+\cdots+k_r \geq 3$ by

$$r_n := \mathrm{Tr}\,\rho\left\{\sum_{k_1+\cdots+k_r\geq 3}\left(\frac{\sqrt{-1}}{\sqrt{n}}\right)^{(k_1+\cdots+k_r)}\prod_{t=1}^r\frac{1}{k_t!}\left(\xi_t^i A_i+\eta_t P(n)\right)^{k_t}\right\}.$$

Then

$$\begin{aligned}
|r_n| &\le \sum_{k_1+\cdots+k_r\ge 3} \left\| \left(\frac{1}{\sqrt{n}}\right)^{(k_1+\cdots+k_r)} \prod_{t=1}^{r} \frac{1}{k_t!}\left(\xi_t^i A_i + \eta_t P(n)\right)^{k_t} \right\| \\
&\le \frac{1}{n\sqrt{n}} \sum_{k_1+\cdots+k_r\ge 3} \left\| \prod_{t=1}^{r} \frac{1}{k_t!}\left(\xi_t^i A_i + \eta_t P(n)\right)^{k_t} \right\| \\
&\le \frac{1}{n\sqrt{n}} \sum_{k_1+\cdots+k_r\ge 3} \prod_{t=1}^{r} \frac{1}{k_t!}\left\|\xi_t^i A_i + \eta_t P(n)\right\|^{k_t} \\
&\le \frac{1}{n\sqrt{n}} \sum_{k_1,\dots,k_r\in\mathbb{Z}_+} \prod_{t=1}^{r} \frac{1}{k_t!}\left\|\xi_t^i A_i + \eta_t P(n)\right\|^{k_t} \\
&= \frac{1}{n\sqrt{n}} \prod_{t=1}^{r} \exp\left\|\xi_t^i A_i + \eta_t P(n)\right\| \\
&\le \frac{1}{n\sqrt{n}} \exp\left(\sum_{t=1}^{r}\left(\left\|\xi_t^i A_i\right\| + \left\|\eta_t P(n)\right\|\right)\right).
\end{aligned}$$

Since $\lim_{n\to\infty} P(n) = 0$, the operators $P(n)$ are uniformly bounded. As a consequence, $\lim_{n\to\infty} n\,|r_n| = 0$, so that $r_n = o\left(\frac{1}{n}\right)$. Thus we conclude that

$$\begin{aligned}
\lim_{n\to\infty} \operatorname{Tr}\rho^{\otimes n}\left(\prod_{t=1}^{r} \mathrm{e}^{\sqrt{-1}\left(\xi_t^i X_i^{(n)} + \eta_t R^{(n)}\right)}\right) &= \lim_{n\to\infty}\left(1 - \frac{1}{2n}\sum_{t=1}^{r}\xi_t^i\xi_t^j J_{ji} - \frac{1}{n}\sum_{t=1}^{r}\sum_{s=t+1}^{r}\xi_t^i\xi_s^j J_{ji} + o\left(\frac{1}{n}\right)\right)^n \\
&= \exp\left(-\frac{1}{2}\sum_{t=1}^{r}\xi_t^i\xi_t^j J_{ji} - \sum_{t=1}^{r}\sum_{s=t+1}^{r}\xi_t^i\xi_s^j J_{ji}\right) \\
&= \phi\left(\prod_{t=1}^{r}\mathrm{e}^{\sqrt{-1}\xi_t^i X_i}\right).
\end{aligned}$$

The last equation is due to (1) with $h = 0$.

3.2 Proof of Theorem 9

Let $X_1, \dots, X_r, \Delta_1, \dots, \Delta_d$ be the basic canonical observables of CCR $\left(\operatorname{Im}\begin{pmatrix}\Sigma & \tau \\ \tau^* & J\end{pmatrix}\right)$, and $\tilde{\phi}$ the quantum Gaussian state $N\left(\begin{pmatrix}0\\0\end{pmatrix}, \begin{pmatrix}\Sigma & \tau \\ \tau^* & J\end{pmatrix}\right)$ on that CCR. Assumption (5) guarantees that the quantities

$$R_h^{(n)} := \mathcal{L}_h^{(n)} - \left\{h^i\Delta_i^{(n)} - \frac{1}{2}J_{ij}h^ih^jI^{(n)}\right\}$$

enjoy $R_h^{(n)} = o\left(\begin{pmatrix} X^{(n)} \\ \Delta^{(n)} \end{pmatrix}, \rho_{\theta_0}^{(n)}\right)$ for each $h \in \mathbb{R}^d$. Consequently, for a finite subset $\{\xi_t\}_{t=1}^r$ of $\mathbb{C}^d$,

$$\begin{aligned}
&\operatorname{Tr} \rho_{\theta_0+h/\sqrt{n}}^{(n)} \left(\prod_{t=1}^r \mathrm{e}^{\sqrt{-1}\xi_t^i X_i^{(n)}}\right) \\
&\quad = \operatorname{Tr} \left(\mathrm{e}^{\frac{1}{2}\mathcal{L}_h^{(n)}} \rho_{\theta_0}^{(n)} \mathrm{e}^{\frac{1}{2}\mathcal{L}_h^{(n)}}\right) \left(\prod_{t=1}^r \mathrm{e}^{\sqrt{-1}\xi_t^i X_i^{(n)}}\right) \\
&\quad = \mathrm{e}^{-\frac{1}{2}h^i h^j J_{ij}} \operatorname{Tr} \rho_{\theta_0}^{(n)} \mathrm{e}^{\frac{1}{2}\left(h^i \Delta_i^{(n)} + R_h^{(n)}\right)} \left(\prod_{t=1}^r \mathrm{e}^{\sqrt{-1}\xi_t^i X_i^{(n)}}\right) \mathrm{e}^{\frac{1}{2}\left(h^i \Delta_i^{(n)} + R_h^{(n)}\right)} \\
&\quad = \mathrm{e}^{-\frac{1}{2}h^i h^j J_{ij}} \operatorname{Tr} \rho_{\theta_0}^{(n)} \left(\mathrm{e}^{-\sqrt{-1}\left(\frac{\sqrt{-1}}{2} h^i \Delta_i^{(n)} + \frac{\sqrt{-1}}{2} R_h^{(n)}\right)}\right) \left(\prod_{t=1}^r \mathrm{e}^{\sqrt{-1}\xi_t^i X_i^{(n)}}\right) \left(\mathrm{e}^{-\sqrt{-1}\left(\frac{\sqrt{-1}}{2} h^i \Delta_i^{(n)} + \frac{\sqrt{-1}}{2} R_h^{(n)}\right)}\right).
\end{aligned}$$

Since $R_h^{(n)}$ is infinitesimal relative to the convergence (4), we see from (2) that

$$\begin{aligned}
&\lim_{n\to\infty} \operatorname{Tr} \rho_{\theta_0+h/\sqrt{n}}^{(n)} \left(\prod_{t=1}^r \mathrm{e}^{\sqrt{-1}\xi_t^i X_i^{(n)}}\right) \\
&\quad = \mathrm{e}^{-\frac{1}{2}h^i h^j J_{ij}} \tilde{\phi} \left(\mathrm{e}^{-\sqrt{-1}\frac{\sqrt{-1}}{2} h^i \Delta_i} \left(\prod_{t=1}^r \mathrm{e}^{\sqrt{-1}\xi_t^i X_i}\right) \mathrm{e}^{-\sqrt{-1}\frac{\sqrt{-1}}{2} h^i \Delta_i}\right) \\
&\quad = \mathrm{e}^{-\frac{1}{2}h^i h^j J_{ij}} \exp\left(-\frac{1}{2}\sum_{t=0}^{r+1} \tilde{\xi}_t^i \tilde{\xi}_t^j \tilde{\Sigma}_{ji} - \sum_{t=0}^{r+1}\sum_{s=t+1}^{r+1} \tilde{\xi}_t^i \tilde{\xi}_s^j \tilde{\Sigma}_{ji}\right) \\
&\quad = \mathrm{e}^{-\frac{1}{2}h^i h^j J_{ij}} \exp\left(-\frac{1}{2}\left\{-\frac{1}{4}h^j h^j J_{ji} + \sum_{t=1}^r \xi_t^i \xi_t^j \Sigma_{ji} - \frac{1}{4}h^j h^j J_{ji}\right\}\right) \\
&\qquad\quad \times \exp\left(\frac{\sqrt{-1}}{2}\sum_{t=1}^r h^i \xi_t^j \tau_{ji} + \frac{\sqrt{-1}}{2}\sum_{t=1}^r \xi_t^i h^j \tau_{ji} + \frac{1}{4}h^i h^j J_{ji} - \sum_{t=1}^r \sum_{s=t+1}^r \xi_t^i \xi_s^j \Sigma_{ji}\right) \\
&\quad = \exp\left(\sum_{t=1}^r \left(\sqrt{-1}\xi_t^i h^j \left(\operatorname{Re}\tau\right)_{ij} - \frac{1}{2}\xi_t^i \xi_t^j \Sigma_{ji}\right) - \sum_{t=1}^r \sum_{s=t+1}^r \xi_t^i \xi_s^j \Sigma_{ji}\right),
\end{aligned}$$

where $\tilde{\Sigma} := \begin{pmatrix} \Sigma & \tau \\ \tau^* & J \end{pmatrix}$ and $(\tilde{\xi}_0, \tilde{\xi}_1, \ldots, \tilde{\xi}_r, \tilde{\xi}_{r+1}) := (-\frac{\sqrt{-1}}{2}h, \xi_1, \ldots, \xi_r, -\frac{\sqrt{-1}}{2}h)$, and (1) was used at the second equation. This is the quasi-characteristic function of $N((\operatorname{Re}\tau)\, h, \Sigma)$.

3.3 Proof of Theorem 10

Since

$$\begin{aligned}
\rho_\theta^{\otimes n} &= \left(\mathrm{e}^{\frac{1}{2}\mathcal{L}\left(\rho_\theta|\rho_{\theta_0}\right)} \rho_{\theta_0} \mathrm{e}^{\frac{1}{2}\mathcal{L}\left(\rho_\theta|\rho_{\theta_0}\right)}\right)^{\otimes n} \\
&= \left(\mathrm{e}^{\frac{1}{2}\sum_{k=1}^n I^{\otimes(k-1)}\otimes\mathcal{L}\left(\rho_\theta|\rho_{\theta_0}\right)\otimes I^{\otimes(n-k)}}\right) \rho_{\theta_0}^{\otimes n} \left(\mathrm{e}^{\frac{1}{2}\sum_{k=1}^n I^{\otimes(k-1)}\otimes\mathcal{L}\left(\rho_\theta|\rho_{\theta_0}\right)\otimes I^{\otimes(n-k)}}\right),
\end{aligned}$$

we see that

$$\mathcal{L}\left(\rho_\theta^{\otimes n} \left|\rho_{\theta_0}^{\otimes n}\right.\right) = \sum_{k=1}^n I^{\otimes(k-1)} \otimes \mathcal{L}\left(\rho_\theta|\rho_{\theta_0}\right) \otimes I^{\otimes(n-k)}. \tag{6}$$

This proves $\rho_\theta^{\otimes n} \sim \rho_{\theta_0}^{\otimes n}$ for all $\theta \in \Theta$ and $n \in \mathbb{N}$.

Before proceeding to the proof of (ii) and (iii) in Definition 8, we give some preliminary consideration. Let the quantum log-likelihood ratio $\mathcal{L}_h := \mathcal{L}\left(\rho_{\theta_0+h}|\rho_{\theta_0}\right)$ be expanded into

$$\mathcal{L}_h = h^i A_i + B_{ij}h^i h^j + o(h^2),$$

where A_i $(1 \le i \le d)$ and B_{ij} $(1 \le i, j \le d)$ are Hermitian operators on $\mathcal{H}$. Observe that A_i is the SLD in the ith direction. In fact,

$$\begin{aligned} \rho_{\theta_0+h} &= \exp\left[\frac{1}{2}\left(h^i A_i + o(h)\right)\right] \rho_{\theta_0} \exp\left[\frac{1}{2}\left(h^i A_i + o(h)\right)\right] \\ &= \rho_{\theta_0} + \frac{1}{2} h^i \left(\rho_{\theta_0} A_i + A_i \rho_{\theta_0}\right) + o(h), \end{aligned}$$

so that

$$\partial_i \rho_{\theta_0} = \frac{1}{2}\left(\rho_{\theta_0} A_i + A_i \rho_{\theta_0}\right).$$

This observation also shows that $\mathrm{Tr}\, \rho_{\theta_0} A_i = 0$ for all i. On the other hand,

$$\begin{aligned} \mathrm{Tr}\, \rho_{\theta_0+h} &= \mathrm{Tr}\, \rho_{\theta_0} \exp\left(h^i A_i + B_{ij} h^i h^j + o(h^2)\right) \\ &= \mathrm{Tr}\, \rho_{\theta_0} \left(I + \left(h^i A_i + B_{ij} h^i h^j\right) + \frac{1}{2}\left(h^i A_i + B_{ij} h^i h^j\right)^2 + o(h^2)\right) \\ &= 1 + h^i \left(\mathrm{Tr}\, \rho_{\theta_0} A_i\right) + h^i h^j \mathrm{Tr}\, \rho_{\theta_0} \left(B_{ij} + \frac{1}{2} A_i A_j\right) + o(h^2) \\ &= 1 + h^i h^j \mathrm{Tr}\, \rho_{\theta_0} \left(B_{ij} + \frac{1}{2} A_i A_j\right) + o(h^2). \end{aligned}$$

Since $\mathrm{Tr}\, \rho_{\theta_0+h} = 1$ for all h, the above equation leads to

$$\mathrm{Tr}\, \rho_{\theta_0} \left(B_{ij} + \frac{1}{2} A_i A_j\right) = 0. \tag{7}$$

Now we prove (ii). Let $J_{ij} := \mathrm{Tr}\, \rho_{\theta_0} A_j A_i$, and let

$$\Delta_i^{(n)} := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes A_i \otimes I^{\otimes(n-k)}.$$

It then follows from the quantum central limit theorem (Proposition 5) that $\left(\Delta^{(n)}, \rho_{\theta_0}^{\otimes n}\right) \underset{q}{\rightsquigarrow} N(0, J)$.

Finally, we prove (iii). It follows from (6) that

$$\mathcal{L}_h^{(n)} = \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes \mathcal{L}_{h/\sqrt{n}} \otimes I^{\otimes(n-k)}.$$

Let us show that

$$R_h^{(n)} := \mathcal{L}_h^{(n)} - \left(h^i \Delta_i^{(n)} - \frac{1}{2}\left(J_{ij} h^i h^j\right) I^{\otimes n}\right)$$

is infinitesimal relative to the convergence $\left(\Delta^{(n)}, \rho_{\theta_0}^{\otimes n}\right) \underset{q}{\rightsquigarrow} N(0, J)$. It is rewritten as

$$\begin{aligned} R_h^{(n)} &= \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes \left[\mathcal{L}_{h/\sqrt{n}} - \frac{1}{\sqrt{n}} h^i A_i + \frac{1}{2n}\left(J_{ij} h^i h^j\right) I\right] \otimes I^{\otimes(n-k)} \\ &= \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes \left[\frac{1}{\sqrt{n}} h^i A_i + \frac{1}{n} B_{ij} h^i h^j + o(\frac{1}{n}) - \frac{1}{\sqrt{n}} h^i A_i + \frac{1}{2n}\left(J_{ij} h^i h^j\right) I\right] \otimes I^{\otimes(n-k)} \\ &= \sum_{k=1}^{n} I^{\otimes(k-1)} \otimes \left[\frac{1}{n} B_{ij} h^i h^j + \frac{1}{2n}\left(J_{ij} h^i h^j\right) I + o(\frac{1}{n})\right] \otimes I^{\otimes(n-k)} \\ &= \sum_{k=1}^{n} I^{\otimes k-1} \otimes \frac{1}{\sqrt{n}} P(n) \otimes I^{\otimes(n-k)}, \end{aligned}$$

where

$$P(n) := \sqrt{n}\left(\frac{1}{n}\left(B_{ij} + \frac{1}{2}J_{ij}I\right)h^i h^j + o(\frac{1}{n})\right).$$

Note that $\lim_{n\to\infty} P(n) = 0$, and that

$$\begin{aligned}
\lim_{n\to\infty}\sqrt{n}\,\mathrm{Tr}\,\rho_{\theta_0}P(n) &= \mathrm{Tr}\,\rho_{\theta_0}\left(B_{ij} + \frac{1}{2}J_{ij}I\right)h^i h^j \\
&= \mathrm{Tr}\,\rho_{\theta_0}\left(B_{ij} + \frac{1}{2}J_{ji}I\right)h^i h^j \\
&= \mathrm{Tr}\,\rho_{\theta_0}\left(B_{ij} + \frac{1}{2}A_i A_j\right)h^i h^j \\
&= 0
\end{aligned}$$

because of (7). It then follows from Lemma 6 that $R_h^{(n)} = o(\Delta^{(n)}, \rho_{\theta_0}^{\otimes n})$ for each $h \in \mathbb{R}^d$. This completes the proof.

3.4 Proof of Corollary 11

That $\rho_\theta^{\otimes n} \sim \rho_{\theta_0}^{\otimes n}$ was proven in the proof of Theorem 10. Let $\Delta_1^{(n)}, \dots, \Delta_d^{(n)}$ be as in the proof of Theorem 10. It then follows from the quantum central limit theorem that

$$\left(\begin{pmatrix} X^{(n)} \\ \Delta^{(n)} \end{pmatrix}, \rho_{\theta_0}^{\otimes n}\right) \underset{q}{\rightsquigarrow} N\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \Sigma & \tau \\ \tau^* & J \end{pmatrix}\right). \tag{8}$$

Further, because of Lemma 6, the sequence $R_h^{(n)}$ of observables given in the proof of Theorem 10 is also infinitesimal relative to the convergence (8). Now that $(\rho_\theta^{\otimes n}, X^{(n)})$ are jointly QLAN at θ_0, the property $\left(X^{(n)}, \rho_{\theta_0+h/\sqrt{n}}^{\otimes n}\right) \underset{q}{\rightsquigarrow} N((\mathrm{Re}\,\tau)h, \Sigma)$ is an immediate consequence of Theorem 9. This completes the proof.

4 Applications to quantum statistics

Quantum Le Cam's third lemma (Corollary 11) implies convergence of $\left\{\rho_{\theta_0+h/\sqrt{n}}^{\otimes n}\,;\, h \in \mathbb{R}^d\right\}$ to a quantum Gaussian shift model $\left\{N((\mathrm{Re}\,\tau)\,h,\,\Sigma)\,;\, h \in \mathbb{R}^d\right\}$. This fact prompts us to expect that, for sufficiently large n, the estimation problem for the parameter h of $\rho_{\theta_0+h/\sqrt{n}}^{\otimes n}$ could be reduced to that for the shift parameter h of a quantum Gaussian shift model $N((\mathrm{Re}\,\tau)\,h,\,\Sigma)$. The latter problem has been well-established to date (see Appendix B). In particular, the best strategy for estimating the shift parameter h is the one that achieves the Holevo bound $C_h\left(N((\mathrm{Re}\,\tau)\,h, \Sigma),\, G\right)$, (see Theorem 25). Moreover, it can be shown (see Corollary 24) that the Holevo bound $C_h\left(N((\mathrm{Re}\,\tau)\,h, \Sigma),\, G\right)$ is identical to the Holevo bound $C_{\theta_0}\left(\rho_\theta,\, G\right)$ for the model ρ_θ at θ_0 This observation strongly suggests the existence of a sequence $M^{(n)}$ of estimators for the parameter h of $\left\{\rho_{\theta_0+h/\sqrt{n}}^{\otimes n}\right\}_n$ that asymptotically achieves the Holevo bound $C_{\theta_0}\left(\rho_\theta,\, G\right)$.

This section is devoted to materialize this program: we prove that there exist a sequence $M^{(n)}$ of estimators on $\left\{\rho_{\theta_0+h/\sqrt{n}}^{\otimes n}\right\}_n$ that is asymptotically unbiased and achieves the Holevo bound $C_{\theta_0}\left(\rho_\theta, G\right)$ for all h that belong to a dense subset of $\mathbb{R}^d$. Since this result requires only twice differentiability of the quantum log-likelihood ratio of the model ρ_θ, it will be useful in a wide range of statistical estimation problems.

4.1 Achievability of the Holevo bound

Theorem 12. *Let* $\left\{\rho_\theta\,;\,\theta\in\Theta\subset\mathbb{R}^d\right\}$ *be a quantum statistical model on a finite dimensional Hilbert space* $\mathcal{H}$*, and fix a point* $\theta_0\in\Theta$*. Suppose that* $\rho_\theta\sim\rho_{\theta_0}$ *for all* $\theta\in\Theta$*, and that the quantum log-likelihood ratio* $\mathcal{L}_h:=\mathcal{L}\left(\rho_{\theta_0+h}|\rho_{\theta_0}\right)$ *is differentiable in* h *around* $h=0$ *and twice differentiable at* $h=0$*. For any countable dense subset* D *of* $\mathbb{R}^d$ *and any weight matrix* G*, there exist a sequence* $M^{(n)}$ *of estimators on the model* $\left\{\rho_{\theta_0+h/\sqrt{n}}^{\otimes n}\,;\,h\in\mathbb{R}^d\right\}$ *that enjoys*

$$\lim_{n\to\infty}E_h^{(n)}[M^{(n)}]=h$$

and

$$\lim_{n\to\infty}\operatorname{Tr}GV_h^{(n)}[M^{(n)}]=C_{\theta_0}\left(\rho_\theta,G\right)$$

for every $h\in D$*. Here* $C_{\theta_0}\left(\rho_\theta,G\right)$ *is the Holevo bound at* θ_0*.*

Proof. Let $\mathcal{D}:=\mathcal{D}_{\rho_{\theta_0}}$ be the commutation operator with respect to the state ρ_{θ_0} (see Appendix A), and let $\mathcal{T}$ be the minimal $\mathcal{D}$ invariant extension of the SLD tangent space $\operatorname{span}_{\mathbb{R}}\left\{L_i\right\}_{i=1}^d$ of the model $\{\rho_\theta\}$ at $\theta=\theta_0$, i.e., the smallest $\mathcal{D}$ invariant real linear subspace of Hermitian operators on $\mathcal{H}$ containing all the SLDs $\{L_i\}_{i=1}^d$ of ρ_θ at θ_0. The minimality ensures that $\operatorname{Tr}\rho_{\theta_0}A=0$ for all $A\in\mathcal{T}$ because $\mathcal{T}'=\{A\in\mathcal{T}\,;\,\operatorname{Tr}\rho_{\theta_0}A=0\}$ is also $\mathcal{D}$ invariant.

Let $\{D_j\}_{j=1}^r$ be a basis of $\mathcal{T}$, thus $d\le r$. Let Σ be an $r\times r$ matrix whose (i,j)th entry is given by $\Sigma_{ij}=\operatorname{Tr}\rho_{\theta_0}D_jD_i$, and let τ be an $r\times d$ matrix whose (i,j)th entry is given by $\tau_{ij}=\operatorname{Tr}\rho_{\theta_0}L_jD_i$. According to Theorem 19 in Appendix A, the Holevo bound for a weight $G>0$ can be expressed as

$$\begin{aligned}C_{\theta_0}\left(\rho_\theta,G\right) &= \min_F\{\operatorname{Tr}GZ+\operatorname{Tr}\left|\sqrt{G}\operatorname{Im}Z\sqrt{G}\right|\,;\,Z={}^tF\Sigma F,\\ &\qquad F\text{ is an }r\times d\text{ real matrix satisfying }{}^tF\operatorname{Re}(\tau)=I\}.\end{aligned}\tag{9}$$

Letting

$$X_i^{(n)}:=\frac{1}{\sqrt{n}}\sum_{k=1}^nI^{\otimes(k-1)}\otimes D_i\otimes I^{\otimes(n-k)}\quad(1\le i\le r),$$

Corollary 11 asserts that $\left(\left\{\rho_\theta^{\otimes n}\right\},X^{(n)}\right)$ is jointly QLAN at θ_0, and that

$$\left(X^{(n)},\rho_{\theta_0+h/\sqrt{n}}^{\otimes n}\right)\underset{q}{\rightsquigarrow}N((\operatorname{Re}\tau)h,\Sigma).\tag{10}$$

Let F be the matrix that attains the minimum in (9), and let $Z:={}^tF\Sigma F$, $\tilde{V}:=\operatorname{Re}Z$, $\tilde{S}:=\operatorname{Im}Z$, $\hat{V}=\sqrt{G^{-1}}\left|\sqrt{G}\operatorname{Im}Z\sqrt{G}\right|\sqrt{G^{-1}}$, and $\hat{Z}=\hat{V}-\sqrt{-1}\tilde{S}$. It then follows from Corollary 24 and Theorem 25 in Appendix B that

$$C_{\theta_0}\left(\rho_\theta,G\right)=\operatorname{Tr}G\left(\tilde{V}+\hat{V}\right).$$

Further, Lemma 13 below assures that there exist a finite dimensional Hilbert space $\hat{\mathcal{H}}$ and a state σ and observables B_i $(1\le i\le d)$ on $\hat{\mathcal{H}}$ such that $\operatorname{Tr}\sigma B_i=0$ and $\operatorname{Tr}\sigma B_jB_i=\hat{Z}_{ij}$. Let

$$\overline{X}_i^{(n)}:=\tilde{X}_i^{(n)}\otimes\hat{I}^{\otimes n}+I^{\otimes n}\otimes Y_i^{(n)}\quad(1\le i\le d),$$

where $\tilde{X}^{(n)}:=F_i^kX_k^{(n)}$ $(1\le i\le d)$,

$$Y_i^{(n)}:=\frac{1}{\sqrt{n}}\sum_{k=1}^n\hat{I}^{\otimes(k-1)}\otimes B_i\otimes\hat{I}^{\otimes(n-k)}\quad(1\le i\le d),$$

and $\hat{I}$ is the identity on $\hat{\mathcal{H}}$. A crucial observation is that $\left(\overline{X}^{(n)}, \overline{\rho}_h^{(n)}\right)$, where $\overline{\rho}_h^{(n)} := \rho_{\theta_0+h/\sqrt{n}}^{\otimes n} \otimes \sigma^{\otimes n}$, converges to a classical Gaussian state:

$$\left(\overline{X}^{(n)}, \overline{\rho}_h^{(n)}\right) \underset{q}{\rightsquigarrow} N(h, \tilde{V} + \hat{V}), \tag{11}$$

for all $h \in \mathbb{R}^d$. In fact,

$$\begin{aligned}
\lim_{n\to\infty} \operatorname{Tr} \overline{\rho}_h^{(n)} \left(\prod_{t=1}^{s} \mathrm{e}^{\sqrt{-1}\xi_t^i \overline{X}_i^{(n)}} \right) &= \lim_{n\to\infty} \operatorname{Tr} \overline{\rho}_h^{(n)} \left\{ \left(\prod_{t=1}^{s} \mathrm{e}^{\sqrt{-1}\xi_t^i \tilde{X}_i^{(n)}} \right) \otimes \left(\prod_{t=1}^{s} \mathrm{e}^{\sqrt{-1}\xi_t^i Y_i^{(n)}} \right) \right\} \\
&= \lim_{n\to\infty} \left[\operatorname{Tr} \rho_{\theta_0+h/\sqrt{n}}^{\otimes n} \left(\prod_{t=1}^{s} \mathrm{e}^{\sqrt{-1}\xi_t^i \tilde{X}_i^{(n)}} \right) \right] \left[\operatorname{Tr} \sigma^{\otimes n} \left(\prod_{t=1}^{s} \mathrm{e}^{\sqrt{-1}\xi_t^i Y_i^{(n)}} \right) \right] \\
&= \phi_h \left(\prod_{t=1}^{s} \mathrm{e}^{\sqrt{-1}\xi_t^i \tilde{X}_i} \right) \psi \left(\prod_{t=1}^{s} \mathrm{e}^{\sqrt{-1}\xi_t^i Y_i} \right),
\end{aligned} \tag{12}$$

where $\tilde{X}_i := F_i^k X_k$ $(1 \le i \le d)$ are canonical observables with $X_1, \ldots, X_r$ being the basic canonical observables of $\mathrm{CCR}\,(\operatorname{Im}\Sigma)$ and $(X, \phi_h) \sim N((\operatorname{Re}\tau)h, \Sigma)$, and $Y_1, \ldots, Y_d$ are the basic canonical observables of $\mathrm{CCR}\left(\operatorname{Im}\hat{Z}\right)$ with $(Y, \psi) \sim N(0, \hat{Z})$. In the last line in (12), we used (10) as well as the quantum central limit theorem for $Y^{(n)}$. By using the explicit form (1) of the quasi-characteristic function for the quantum Gaussian state, (12) is rewritten as

$$\begin{aligned}
&\exp\left(\sum_{t=1}^{r} \left(\sqrt{-1}\xi_t^i h_i - \frac{1}{2}\xi_t^i \xi_t^j Z_{ji} \right) - \sum_{t=1}^{r} \sum_{s=t+1}^{r} \xi_t^i \xi_s^j Z_{ji} \right) \exp\left(-\frac{1}{2} \sum_{t=1}^{r} \xi_t^i \xi_t^j \hat{Z}_{ji} - \sum_{t=1}^{r} \sum_{s=t+1}^{r} \xi_t^i \xi_s^j \hat{Z}_{ji} \right) \\
&\qquad = \exp\left(\sum_{t=1}^{r} \left(\sqrt{-1}\xi_t^i h_i - \frac{1}{2}\xi_t^i \xi_t^j (\tilde{V} + \hat{V})_{ji} \right) - \sum_{t=1}^{r} \sum_{s=t+1}^{r} \xi_t^i \xi_s^j (\tilde{V} + \hat{V})_{ji} \right).
\end{aligned}$$

This proves (11).

Now according to Lemma 14 below, there exist a quintuple sequence

$$M^{(n,m,\ell,q,p)} = \left\{ M_\omega^{(n,m,\ell,q,p)} \,;\ \omega \in \Omega^{(n,m,l,p,q)} \right\}$$

of POVMs on $\left(\mathcal{H} \otimes \hat{\mathcal{H}}\right)^{\otimes n}$, taking values in a certain finite subset $\Omega^{(n,m,l,p,q)}$ of $\mathbb{R}^d$, that enjoys the properties

$$\lim_{p\to\infty} \lim_{q\to\infty} \lim_{\ell\to\infty} \lim_{m\to\infty} \lim_{n\to\infty} \overline{E}_h^{(n)}[M^{(n,m,\ell,q,p)}] = h,$$

and

$$\lim_{p\to\infty} \lim_{q\to\infty} \lim_{\ell\to\infty} \lim_{m\to\infty} \lim_{n\to\infty} \overline{V}_h^{(n)}[M^{(n,m,\ell,q,p)}] = \tilde{V} + \hat{V},$$

for all $h \in \mathbb{R}^d$, where $\overline{E}_h^{(n)}[\,\cdot\,]$ and $\overline{V}_h^{(n)}[\,\cdot\,]$ denote the expectation and the covariance with respect to $\overline{\rho}_h^{(n)}$. It then follows from Lemma 15 below that for any countable dense subset D of $\mathbb{R}^d$ and any $h \in D$, there exist a subsequence $\{(n, m(n), \ell(n), q(n), p(n)\}_{n\in\mathbb{N}}$ such that

$$\lim_{n\to\infty} \overline{E}_h^{(n)}[M^{(n,m(n),\ell(n),q(n),p(n))}] = h,$$

and

$$\lim_{n\to\infty} \overline{V}_h^{(n)}[M^{(n,m(n),\ell(n),q(n),p(n))}] = \tilde{V} + \hat{V}.$$

This implies that the POVM $M^{(n)}$ on $\mathcal{H}^{\otimes n}$ that is uniquely defined by the requirement

$$\operatorname{Tr} \rho^{(n)} M_\omega^{(n)} = \operatorname{Tr} \left(\rho^{(n)} \otimes \sigma^{\otimes n} \right) M_\omega^{(n,m(n),\ell(n),q(n),p(n))}$$

for all density operator $\rho^{(n)}$ on $\mathcal{H}^{\otimes n}$ and $\omega \in \Omega^{(n,m(n),l(n),p(n),q(n))}$ enjoys

$$\lim_{n\to\infty} E_h^{(n)}[M^{(n)}] = h,$$

$$\lim_{n\to\infty} V_h^{(n)}[M^{(n)}] = \tilde{V} + \hat{V}.$$

for all $h \in D$. Recalling that $\operatorname{Tr} G(\tilde{V}+\hat{V}) = C_{\theta_0}(\rho_\theta, G)$, the proof is complete. □

Lemma 13. *Given a $d\times d$ positive semidefinite Hermitian matrix J, there exist a finite dimensional Hilbert space $\mathcal{H}$ and a pure state ρ and observables A_i $(1 \le i \le d)$ on $\mathcal{H}$ such that $\operatorname{Tr} \rho A_i = 0$ and $\operatorname{Tr} \rho A_j A_i = J_{ij}$.*

Proof. Let $\mathcal{H} = \mathbb{C}^{d+1}$, and let $\{|i\rangle\}_{i=0}^d$ be a CONS of $\mathcal{H}$. We set $|\psi\rangle := |0\rangle$ and $|\ell_i\rangle := \sum_{k=1}^d \left[\sqrt{J}\right]_{ik} |k\rangle$ for $i = 1, \ldots, d$. Then $\rho := |\psi\rangle\langle\psi|$ and $A_i := |\ell_i\rangle\langle\psi| + |\psi\rangle\langle\ell_i|$ satisfy $\operatorname{Tr}\rho A_i = 0$ and $\operatorname{Tr}\rho A_j A_i = J_{ij}$. □

Lemma 14. *Given a sequence $\mathcal{H}^{(n)}$ of finite dimensional Hilbert spaces, let $X^{(n)} = \left(X_1^{(n)}, \ldots, X_d^{(n)}\right)$ be a list of observables on $\mathcal{H}^{(n)}$, and let $\left\{\rho_h^{(n)}\right\}_h$ be a family of density operators on $\mathcal{H}^{(n)}$ parametrized by $h \in \mathbb{R}^d$. If there is a real $d \times d$ positive definite matrix V such that*

$$\left(X^{(n)}, \rho_h^{(n)}\right) \underset{q}{\rightsquigarrow} N(h, V) \tag{13}$$

holds for all $h \in \mathbb{R}^d$, then there exist a quintuple sequence $\left\{M^{(n,m,\ell,q,p)}; (n,m,\ell,q,p) \in \mathbb{N}^5\right\}$ of POVMs on $\mathcal{H}^{(n)}$ that enjoy the properties

$$\lim_{p\to\infty}\lim_{q\to\infty}\lim_{\ell\to\infty}\lim_{m\to\infty}\lim_{n\to\infty} E_h^{(n)}[M^{(n,m,\ell,q,p)}] = h,$$

and

$$\lim_{p\to\infty}\lim_{q\to\infty}\lim_{\ell\to\infty}\lim_{m\to\infty}\lim_{n\to\infty} V_h^{(n)}[M^{(n,m,\ell,q,p)}] = V.$$

Proof. Let

$$\Omega^{(m,\ell)} := \left\{ \frac{\ell}{m}\overrightarrow{k} + \frac{\ell}{2m}(1, \ldots, 1)\,;\ \overrightarrow{k} \in \mathbb{Z}^d \right\} \cap [-l, l]^d$$

be a finite subset of $\mathbb{R}^d$, comprising $(2m)^d$ lattice points in the hypercube $[-l,l]^d$, and let $\Omega^{(m,\ell,p)} := \Omega^{(m,\ell)} \cap [-p,p]^d$ and $\Omega_0^{(m,\ell,p)} := \Omega^{(m,\ell,p)} \cup \{0\}$. We introduce a Gaussian density function $f_\omega^{(q)}(x)$ on $\mathbb{R}^d$ centered at $\omega = (\omega_1, \ldots, \omega_d) \in \mathbb{R}^d$ by

$$f_\omega^{(q)}(x) := \left\{\prod_{i=1}^d g_{\omega_{d+1-i}}^{(q)}(x_{d+1-i})\right\}\left\{\prod_{i=1}^d g_{\omega_i}^{(q)}(x_i)\right\},$$

where $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$ and

$$g_s^{(q)}(t) := \left(\frac{q}{2\pi}\right)^{\frac{1}{4}} \exp\left(-\frac{q}{4}(t-s)^2\right), \qquad (s, t \in \mathbb{R}).$$

By using this function, we define a POVM $M^{(n,m,l,q,p)} = \left\{M_\omega^{(n,m,l,q,p)}; \omega \in \Omega_0^{(m,\ell,p)}\right\}$ on $\mathcal{H}^{(n)}$ that takes values in the finite subset $\Omega_0^{(m,\ell,p)}$ by

$$M_\omega^{(n,m,\ell,q,p)} := R^{(m,\ell,q)}(X^{(n)}) \left[f_\omega^{(q)}(X^{(n)}) + \frac{I^{(n)}}{(2m)^d}\right] R^{(m,\ell,q)}(X^{(n)})$$

for $\omega \in \Omega^{(m,\ell,p)}$, and

$$M_0^{(n,m,\ell,q,p)} := \sum_{\omega\in\Omega^{(m,\ell)}\setminus\Omega^{(m,\ell,p)}} \left\{ R^{(m,\ell,q)}(X^{(n)}) \left[\left(f_\omega^{(q)}(X^{(n)}) + \frac{I^{(n)}}{(2m)^d} \right) \right] R^{(m,\ell,q)}(X^{(n)}) \right\}.$$

Here

$$R^{(m,\ell,q)}(x) := g\left(\sum_{\omega\in\Omega^{(m,\ell)}} f_\omega^{(q)}(x) \right)$$

is the normalization with

$$g(t) := \frac{1}{\sqrt{t+1}}.$$

Intuitively speaking, the difference set $\Omega^{(m,\ell)} \setminus \Omega^{(m,\ell,p)}$ works as a "buffer" zone that gives the default outcome $\omega = 0$. This device is meaningful only when $p < \ell$.

We shall prove that

$$\lim_{p\to\infty}\lim_{q\to\infty}\lim_{\ell\to\infty}\lim_{m\to\infty}\lim_{n\to\infty} \sum_{\omega\in\Omega_0^{(m,\ell,p)}} P(\omega)\mathrm{Tr}\,\rho_h^{(n)} M_\omega^{(n,m,\ell,q,p)} = \int_{\mathbb{R}^d} P(\omega)p_h(\omega)\mathrm{d}\omega, \tag{14}$$

where $P(\omega)$ is an arbitrary polynomial of ω such that $P(0) = 0$ and $p_h(\omega)$ is a probability density function of the classical normal distribution $N(h, V)$. Once (14) has been proved, we can verify

$$\lim_{p\to\infty}\lim_{q\to\infty}\lim_{\ell\to\infty}\lim_{m\to\infty}\lim_{n\to\infty} E_h^{(n)}[M^{(n,m,\ell,q,p)}] = h$$

and

$$\lim_{p\to\infty}\lim_{q\to\infty}\lim_{\ell\to\infty}\lim_{m\to\infty}\lim_{n\to\infty} V_h^{(n)}[M^{(n,m,\ell,q,p)}] = V$$

just by letting $P(\omega) = \omega_i$ or $P(\omega) = \omega_i\omega_j$ $(1 \le i, j \le d)$ in (14).

The first limit $n \to \infty$ in (14) yields

$$\begin{aligned}
&\lim_{n\to\infty} \sum_{\omega\in\Omega_0^{(m,\ell,p)}} P(\omega)\mathrm{Tr}\,\rho_h^{(n)} M_\omega^{(n,m,\ell,q,p)} \\
&= \lim_{n\to\infty} \sum_{\omega\in\Omega^{(m,\ell,p)}} P(\omega)\mathrm{Tr}\,\rho_h^{(n)} M_\omega^{(n,m,\ell,q,p)} \\
&= \lim_{n\to\infty} \sum_{\omega\in\Omega^{(m,\ell,p)}} P(\omega)\mathrm{Tr}\,\rho_h^{(n)} R^{(m,\ell,q)}(X^{(n)}) \left[f_\omega^{(q)}(X^{(n)}) + \frac{I^{(n)}}{(2m)^d} \right] R^{(m,\ell,q)}(X^{(n)}) \\
&= \sum_{\omega\in\Omega^{(m,\ell,p)}} P(\omega) E_h \left[R^{(m,\ell,q)}(X)^2 \left(f_\omega^{(q)}(X) + \frac{I}{(2m)^d} \right) \right] \\
&= \int_{\mathbb{R}^d} \frac{\sum_{\omega\in\Omega^{(m,\ell,p)}} P(\omega)\left(f_\omega^{(q)}(x) + \frac{1}{(2m)^d} \right)}{\sum_{\omega\in\Omega^{(m,\ell)}} \left(f_\omega^{(q)}(x) + \frac{1}{(2m)^d} \right)} p_h(x)\mathrm{d}x.
\end{aligned} \tag{15}$$

In the fourth line, we used the assumption (13) and Corollary 29 in Appendix D, as well as the fact that functions $g_s^{(q)}(t)$ on $\mathbb{R}$ and $g(t)$ on $t \ge 0$ are both bounded and continuous. Further, $X = (X_1, \ldots, X_d)$ is a classical random vector that follow the normal distribution $N(h, V)$, and $E_h[\,\cdot\,]$ denotes the expectation with respect to $N(h, V)$. As for the second limit $m \to \infty$, due to

$$\left| \frac{\sum_{\omega\in\Omega^{(m,\ell,p)}} P(\omega)\left(f_\omega^{(q)}(x) + \frac{1}{(2m)^d} \right)}{\sum_{\omega\in\Omega^{(m,\ell)}} \left(f_\omega^{(q)}(x) + \frac{1}{(2m)^d} \right)} \right| \le \max_{\omega\in[-p,p]^d} |P(\omega)| < \infty,$$

the bounded convergence theorem yields

$$
\begin{aligned}
\lim_{m\to\infty}(15) &= \int_{\mathbb{R}^d} \lim_{m\to\infty} \frac{\left(\frac{\ell}{m}\right)^d \sum_{\omega\in\Omega^{(m,\ell,p)}} P(\omega)\left(f_\omega^{(q)}(x)+\frac{1}{(2m)^d}\right)}{\left(\frac{\ell}{m}\right)^d \sum_{\omega\in\Omega^{(m,\ell)}}\left(f_\omega^{(q)}(x)+\frac{1}{(2m)^d}\right)} p_h(x)\mathrm{d}x \\
&= \int_{\mathbb{R}^d} \frac{\int_{\omega\in[-p,p]^d} P(\omega)p^{(q)}(\omega,x)\mathrm{d}\omega}{\int_{\omega\in[-\ell,\ell]^d} p^{(q)}(\omega,x)\mathrm{d}\omega} p_h(x)\mathrm{d}x, \qquad (16)
\end{aligned}
$$

where $p^{(q)}(\omega,x)=\left(\frac{q}{2\pi}\right)^{\frac{d}{2}}\exp\left(-\frac{q}{2}\sum_{i=1}^d(x_i-\omega_i)^2\right)$, and Darboux's theorem for the Riemann integral was used in the second line. Finally, the dominated convergence theorem and Fubini's theorem yield

$$
\begin{aligned}
\lim_{p\to\infty}\lim_{q\to\infty}\lim_{\ell\to\infty}(16) &= \lim_{p\to\infty}\lim_{q\to\infty}\int_{\mathbb{R}^d} \frac{\int_{\omega\in[-p,p]^d} P(\omega)p^{(q)}(\omega,x)\mathrm{d}\omega}{\int_{\mathbb{R}^d} p^{(q)}(\omega,x)\mathrm{d}\omega} p_h(x)\mathrm{d}x \\
&= \lim_{p\to\infty}\lim_{q\to\infty}\int_{\mathbb{R}^d}\left(\int_{\omega\in[-p,p]^d} P(\omega)p^{(q)}(\omega,x)\mathrm{d}\omega\right)p_h(x)\mathrm{d}x \\
&= \lim_{p\to\infty}\lim_{q\to\infty}\int_{\omega\in[-p,p]^d}\left(\int_{\mathbb{R}^d} p^{(q)}(\omega,x)p_h(x)\mathrm{d}x\right)P(\omega)\mathrm{d}\omega \\
&= \lim_{p\to\infty}\lim_{q\to\infty}\int_{\omega\in[-p,p]^d} p_h^{(q)}(\omega)P(\omega)\mathrm{d}\omega \\
&= \lim_{p\to\infty}\int_{\omega\in[-p,p]^d} p_h(\omega)P(\omega)\mathrm{d}\omega \\
&= \int_{\mathbb{R}^d} p_h(\omega)P(\omega)\mathrm{d}\omega, \qquad (17)
\end{aligned}
$$

where $p_h^{(q)}(\omega)$ is the density function of $N(h, V+\frac{1}{q}I)$. This completes the proof. □

Lemma 15. *For each $i\in\mathbb{N}$, let $\left\{a^i_{n_1n_2\cdots n_rn};\,(n_1,n_2,\ldots,n_r,n)\in\mathbb{N}^{(r+1)}\right\}$ be an $(r+1)$-tuple sequence on a normed space V. If, for each $i\in\mathbb{N}$, there exists an $\alpha^i\in V$ such that*

$$\lim_{n_1\to\infty}\lim_{n_2\to\infty}\cdots\lim_{n_r\to\infty}\lim_{n\to\infty} a^i_{n_1n_2\cdots n_rn}=\alpha^i,$$

then there exist a subsequence $\{(n_1(n),n_2(n),\ldots,n_r(n),n)\}_{n\in\mathbb{N}}$ that satisfies

$$\lim_{n\to\infty} a^i_{n_1(n)n_2(n)\cdots n_r(n)n}=\alpha^i$$

for all $i\in\mathbb{N}$.

Proof. We first prove the case when $r=1$. Let $a^i_{n_1}:=\lim_{n\to\infty}a^i_{n_1n}$. We construct a subsequence $\{(n_1(k),n(k))\}_{k\in\mathbb{N}}$ in a recursive manner as follows. We set $n_1(1)=n(1)=1$. For $k\geq 2$, it follows from $\lim_{n_1\to\infty}a^i_{n_1}=\alpha^i$ that there exist an $N_1(k)\in\mathbb{N}$ such that $n_1\geq N_1(k)$ implies

$$\max_{1\leq i\leq k}\left|a^i_{n_1}-\alpha^i\right|<\frac{1}{k}.$$

Thus the number $n_1(k):=\max\{N_1(k),n_1(k-1)+1\}$ satisfies

$$\max_{1\leq i\leq k}\left|a^i_{n_1(k)}-\alpha^i\right|<\frac{1}{k}. \qquad (18)$$

For this $n_1(k)$, it follows from $\lim_{n\to\infty}a^i_{n_1(k)n}=a^i_{n_1(k)}$ that there exist an $N(k)\in\mathbb{N}$ such that $n\geq N(k)$ implies

$$\max_{1\leq i\leq k}\left|a^i_{n_1(k)n}-a^i_{n_1(k)}\right|<\frac{1}{k}. \qquad (19)$$

Thus we set $n(k) := \max\{N(k), n(k-1)+1\}$.

Now let $k(n) := \max\{k;\, n(k) \le n\}$, which is non-decreasing in n and $\lim_{n\to\infty} k(n) = \infty$. We show that the subsequence $\{n_1(k(n)), n);\, n \in \mathbb{N}\}$ enjoys the required property: for all $i \in \mathbb{N}$,

$$\lim_{n\to\infty} a^i_{n_1(k(n))n} = \alpha^i.$$

Given $i \in \mathbb{N}$ and $\varepsilon > 0$ arbitrarily, there exist an $N \in \mathbb{N}$ such that $n \ge N$ implies $k(n) \ge \max\left\{i, \left\lceil \frac{2}{\varepsilon} \right\rceil\right\}$. Then for all $n \ge N$, we have

$$\begin{aligned}\left|a^i_{n_1(k(n))n} - \alpha^i\right| &\le \left|a^i_{n_1(k(n))n} - a^i_{n_1(k(n))}\right| + \left|a^i_{n_1(k(n))} - \alpha^i\right| \\ &\le \max_{1\le j\le k(n)} \left|a^j_{n_1(k(n))n} - a^j_{n_1(k(n))}\right| + \max_{1\le j\le k(n)} \left|a^j_{n_1(k(n))} - \alpha^j\right| \\ &< \frac{2}{k(n)} \le \varepsilon.\end{aligned}$$

In the third inequality, we used (18) and (19), as well as its premise $n \ge n(k(n)) \ge N(k(n))$.

The proof for a generic r is similar. □

4.2 Application to qubit state estimation

In order to demonstrate the power of our method, we explore qubit state estimation problems.

Example 16 (3-dimensional faithful state model)**.**

The first example is an ordinary one, comprising the totality of faithful quit states:

$$\mathcal{S}(\mathbb{C}^2) = \left\{\rho_\theta = \frac{1}{2}\left(I + \theta^1\sigma_1 + \theta^2\sigma_2 + \theta^3\sigma_3\right)\ ;\ \theta = (\theta^i)_{1\le i\le 3} \in \Theta\right\}$$

where σ_i $(i = 1,2,3)$ are the standard Pauli matrices and Θ is the open unit ball in $\mathbb{R}^3$. Due to the rotational symmetry, we take the reference point to be $\theta_0 = (0,0,r)$, with $0 \le r < 1$. By a direct calculation, we see that the SLDs of the model ρ_θ at $\theta = \theta_0$ are $(L_1, L_2, L_3) = \left(\sigma_1, \sigma_2, (rI+\sigma_3)^{-1}\right)$, and the SLD Fisher information matrix $J^{(S)}$ at θ_0 is given by the real part of the matrix

$$J := \left[\operatorname{Tr} \rho_{\theta_0} L_j L_i\right]_{ij} = \begin{pmatrix} 1 & -r\sqrt{-1} & 0 \\ r\sqrt{-1} & 1 & 0 \\ 0 & 0 & 1/(1-r^2) \end{pmatrix}.$$

Given a 3×3 real positive definite matrix G, the minimal value of the weighted covariances at $\theta = \theta_0$ is given by

$$\min_{\hat{M}} \operatorname{Tr} G V_{\theta_0}[\hat{M}] = C^{(1)}_{\theta_0}(\rho_\theta, G),$$

where the minimum is taken over all estimators $\hat{M}$ that are locally unbiased at θ_0, and

$$C^{(1)}_{\theta_0}(\rho_\theta, G) = \left(\operatorname{Tr}\sqrt{\sqrt{G} J^{(S)^{-1}} \sqrt{G}}\right)^2$$

is the Hayashi-Gill-Massar bound [7, 3] (see also [21]). On the other hand, the SLD tangent space is obviously $\mathcal{D}$ invariant, and the Holevo bound is given by

$$C_{\theta_0}(\rho_\theta, G) := \operatorname{Tr} G J^{(R)^{-1}} + \operatorname{Tr}\left|\sqrt{G} \operatorname{Im} J^{(R)^{-1}} \sqrt{G}\right|,$$

where

$$J^{(R)^{-1}} := (\operatorname{Re} J)^{-1} J (\operatorname{Re} J)^{-1} = \begin{pmatrix} 1 & -r\sqrt{-1} & 0 \\ r\sqrt{-1} & 1 & 0 \\ 0 & 0 & 1-r^2 \end{pmatrix}$$

is the inverse RLD Fisher information matrix (See Corollary 20 in Appendix A).

It can be shown that the Hayashi-Gill-Massar bound is greater than the Holevo bound:

$$C_{\theta_0}^{(1)}(\rho_\theta, G) > C_{\theta_0}(\rho_\theta, G).$$

Let us check this fact for the special case when $G = J^{(S)}$. A direct computation shows that

$$C_{\theta_0}^{(1)}\left(\rho_\theta, J^{(S)}\right) = 9,$$

and

$$C_{\theta_0}\left(\rho_\theta, J^{(S)}\right) = 3 + 2r.$$

The left panel of Figure 1 shows the behavior of $C_{\theta_0}\left(\rho_\theta, J^{(S)}\right)$ (solid) and $C_{\theta_0}^{(1)}\left(\rho_\theta, J^{(S)}\right)$ (dashed) as functions of r. We see that the Holevo bound $C_{\theta_0}\left(\rho_\theta, J^{(S)}\right)$ is much smaller than $C_{\theta_0}^{(1)}\left(\rho_\theta, J^{(S)}\right)$.

Does this fact imply that the Holevo bound is of no use? The answer is contrary, as Theorem 12 asserts. We will demonstrate the asymptotic achievability of the Holevo bound. Let

$$\Delta_i^{(n)} := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} I^{\otimes k-1} \otimes L_i \otimes I^{\otimes n-k}$$

and let $X_i^{(n)} := \Delta_i^{(n)}$ for $i = 1, 2, 3$. It follows from the quantum central limit theorem that

$$\left(\begin{pmatrix} X^{(n)} \\ \Delta^{(n)} \end{pmatrix}, \rho_{\theta_0}^{\otimes n} \right) \underset{q}{\rightsquigarrow} N\left(0, \begin{pmatrix} J & J \\ J & J \end{pmatrix}\right).$$

Since

$$\mathcal{L}(\theta) := \mathcal{L}\left(\rho_\theta | \rho_{\theta_0}\right) = 2 \log \left(\sqrt{\rho_{\theta_0}^{-1}} \sqrt{\sqrt{\rho_{\theta_0}} \rho_\theta \sqrt{\rho_{\theta_0}}} \sqrt{\rho_{\theta_0}^{-1}} \right)$$

is obviously of class C^∞ in θ, Corollary 11 shows that $\left(\left\{\rho_\theta^{\otimes n}\right\}, X^{(n)}\right)$ is jointly QLAN at θ_0, and that

$$\left(X^{(n)}, \rho_{\theta_0 + h/\sqrt{n}}^{\otimes n}\right) \underset{q}{\rightsquigarrow} N((\operatorname{Re} J)h, J)$$

for all $h \in \mathbb{R}^3$. This implies that a sequence of models $\left\{\rho_{\theta_0 + h/\sqrt{n}}^{\otimes n} ; h \in \mathbb{R}^d\right\}$ converges to a quantum Gaussian shift model $\left\{N((\operatorname{Re} J)h,\ J) ; h \in \mathbb{R}^3\right\}$. Note that the imaginary part

$$S = \begin{pmatrix} 0 & -r\sqrt{-1} & 0 \\ r\sqrt{-1} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

of the matrix J determines the $\operatorname{CCR}(S)$, as well as the corresponding basic canonical observables $X = (X^1, X^2, X^3)$. When $r \neq 0$, the above S has the following physical interpretation: X^1 and X^2 form a canonical pair of quantum Gaussian observables, while X^3 is a classical Gaussian random variable. In this way, the matrix J automatically tells us the structure of the limiting quantum Gaussian shift model.



Figure 1: The left panel displays the Holevo bound $C_{(0,0,r)}\left(\rho_\theta, J^{(S)}\right)$ (solid) and the Hayashi-Gill-Massar bound $C^{(1)}_{(0,0,r)}\left(\rho_\theta, J^{(S)}\right)$ (dashed) for the 3-D model $\rho_\theta = \frac{1}{2}\left(I + \theta^1\sigma_1 + \theta^2\sigma_2 + \theta^3\sigma_3\right)$ as functions of $r = \|\theta\|$. The right panel displays the Holevo bound $C_{(0,r)}\left(\rho_\theta, J^{(S)}\right)$ (solid) and the Nagaoka bound $C^{(1)}_{(0,r)}\left(\rho_\theta, J^{(S)}\right)$ (dashed) for the 2-D model $\rho_\theta = \frac{1}{2}\left(I + \theta^1\sigma_1 + \theta^2\sigma_2 + \frac{1}{4}\sqrt{1-\|\theta\|^2}\,\sigma_3\right)$.

Now, the best strategy for estimating the shift parameter h of the quantum Gaussian shift model $\left\{N((\operatorname{Re} J)h,\, J)\,;\, h \in \mathbb{R}^d\right\}$ is the one that achieves the Holevo bound $C_h\left(N((\operatorname{Re} J)\,h, J),\, G\right)$, (cf., Theorem 25 in Appendix B). Moreover, this Holevo bound $C_h\left(N((\operatorname{Re} J)\,h, J),\, G\right)$ is identical to the Holevo bound $C_{\theta_0}\left(\rho_\theta,\, G\right)$ for the model ρ_θ at θ_0, (cf., Corollary 24. Recall that the matrix J is evaluated at θ_0 of the model ρ_θ). Theorem 12 combines these facts, and concludes that there exist a sequence $M^{(n)}$ of estimators on the model $\left\{\rho^{\otimes n}_{\theta_0+h/\sqrt{n}}\,;\, h \in \mathbb{R}^3\right\}$ that is asymptotically unbiased and achieves the common values of the Holevo bound:

$$\lim_{n\to\infty} \operatorname{Tr} G V_h^{(n)}[M^{(n)}] = C_h\left(N((\operatorname{Re} J)h, J), G\right) = C_{\theta_0}\left(\rho_\theta,\, G\right)$$

for all h that belong to a countable dense subset of $\mathbb{R}^3$.

It should be emphasized that the matrix J becomes the identity at the origin $\theta_0 = (0,0,0)$. This means that the limiting Gaussian shift model $\left\{N(h, J)\,;\, h \in \mathbb{R}^3\right\}$ is “classical.” Since such a degenerate case cannot be treated in [5, 11, 8], our method has a clear advantage in applications.

Example 17 (Pure state model)**.**

The second example is to demonstrate that our formulation allows us to treat pure state models. Let us consider the model $\mathcal{S} = \{|\psi(\theta)\rangle\langle\psi(\theta)|\,;\, \theta = (\theta^i)_{1\le i\le 2} \in \Theta\}$ defined by

$$\psi(\theta) := \frac{1}{\sqrt{\cosh\|\theta\|}}\, \mathrm{e}^{\frac{1}{2}\left(\theta^1\sigma_1 + \theta^2\sigma_2\right)} \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

where Θ is an open subset of $\mathbb{R}^2$ containing the origin, and $\|\cdot\|$ denotes the Euclid norm. By a direct computation, the SLDs at $\theta_0 = (0,0)$ are $(L_1,\, L_2) = (\sigma_1,\, \sigma_2)$, and the SLD Fisher information matrix $J^{(S)}$ is the real part of the matrix

$$J = \left[\operatorname{Tr} \rho_{\theta_0} L_j L_i\right]_{ij} = \begin{pmatrix} 1 & -\sqrt{-1} \\ \sqrt{-1} & 1 \end{pmatrix},$$

that is, $J^{(S)} = I$. Since the SLD tangent space is $\mathcal{D}$ invariant [1], the Holevo bound for a weight $G > 0$ is represented as

$$C_{\theta_0}\left(\rho_\theta, G\right) := \operatorname{Tr} G {J^{(R)}}^{-1} + \operatorname{Tr}\left|\sqrt{G}\,\operatorname{Im} {J^{(R)}}^{-1}\sqrt{G}\right|$$

where

$$J^{(R)^{-1}} := (\operatorname{Re} J)^{-1} J (\operatorname{Re} J)^{-1} = \begin{pmatrix} 1 & -\sqrt{-1} \\ \sqrt{-1} & 1 \end{pmatrix}$$

is the inverse RLD Fisher information matrix (see Corollary 20 in Appendix A).

Let us demonstrate that our QLAN is applicable also to pure state models. Let

$$\Delta_i^{(n)} := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} I^{\otimes k-1} \otimes L_i \otimes I^{\otimes n-k}$$

and let $X_i^{(n)} := \Delta_i^{(n)}$ for $i = 1, 2$. It follows from the quantum central limit theorem that

$$\left(\begin{pmatrix} X^{(n)} \\ \Delta^{(n)} \end{pmatrix}, \rho_{\theta_0}^{\otimes n} \right) \underset{q}{\rightsquigarrow} N\left(0, \begin{pmatrix} J & J \\ J & J \end{pmatrix}\right).$$

Since

$$\mathcal{L}(\theta) := \mathcal{L}\left(\rho_\theta | \rho_{\theta_0}\right) = \theta^1 \sigma_1 + \theta^2 \sigma_2 - \log\cosh \|\theta\|$$

is of class C^∞ with respect to θ, it follows from Corollary 11 that $\left(\left\{\rho_\theta^{\otimes n}\right\}, X^{(n)}\right)$ is jointly QLAN at θ_0, and that

$$(X^{(n)}, \rho_{\theta_0 + h/\sqrt{n}}^{\otimes n}) \rightsquigarrow N((\operatorname{Re} J)h, J) = N(h, J^{(R)^{-1}})$$

for all $h \in \mathbb{R}^2$. Theorem 12 further asserts that there exist a sequence $M^{(n)}$ of estimators on the model $\left\{\rho_{\theta_0 + h/\sqrt{n}}^{\otimes n} \,;\, h \in \mathbb{R}^2\right\}$ that is asymptotically unbiased and achieves the Holevo bound:

$$\lim_{n \to \infty} \operatorname{Tr} G V_h^{(n)}[M^{(n)}] = C_h\left(N(h, J^{(R)^{-1}}), G\right) = C_{(0,0)}\left(\rho_\theta, G\right)$$

for all h that belong to a dense subset of $\mathbb{R}^3$. In fact, the sequence $M^{(n)}$ can be taken to be a separable one, making no use of quantum correlations [15]. (See also Appendix C for a simple proof.) Note that the matrix $J^{(R)^{-1}}$ is degenerate, and the derived quantum Gaussian shift model $\left\{N(h, J^{(R)^{-1}})\right\}_h$ is a canonical coherent model [1].

Example 18 (2-dimensional faithful state model)**.**

The third example treats the case when the SLD tangent space is not $\mathcal{D}$ invariant. Let us consider the model

$$\mathcal{S} = \left\{ \rho_\theta = \frac{1}{2}\left(I + \theta^1 \sigma_1 + \theta^2 \sigma_2 + z_0 \sqrt{1 - \|\theta\|^2}\, \sigma_3\right) \,;\, \theta = (\theta^i)_{1 \le i \le 2} \in \Theta \right\},$$

where $0 \le z_0 < 1$, and Θ is the open unit disk. Due to the rotational symmetry around z-axis, we take the reference point to be $\theta_0 = (0, r)$, with $0 \le r < 1$. By a direct calculation, we see that the SLDs at θ_0 are $(L_1, L_2) = \left(\sigma_1, \frac{1}{1-r^2}(\sigma_2 - rI)\right)$. It is important to notice that the SLD tangent space $\operatorname{span}\{L_i\}_{i=1}^2$ is not $\mathcal{D}$ invariant unless $r = 0$. In fact

$$\mathcal{D}\sigma_1 = z(r)\sigma_2 - r\sigma_3, \qquad \mathcal{D}\sigma_2 = -z(r)\sigma_1,$$

where $z(r) := E[\sigma_3] = z_0\sqrt{1 - r^2}$. The minimal $\mathcal{D}$ invariant extension $\mathcal{T}$ of the SLD tangent space has a basis $(D_1, D_2, D_3) := (L_1, L_2, \sigma_3 - z(r)I)$. The matrices Σ, J, and τ appeared in Definition 8 and

Corollary 11 are calculated as

$$\begin{aligned}
\Sigma &:= [\operatorname{Tr}\rho_{\theta_0} D_j D_i]_{ij} = \begin{pmatrix} 1 & -\sqrt{-1}\dfrac{z_0^2}{z(r)} & r\sqrt{-1}-z(r) \\ \sqrt{-1}\dfrac{z_0^2}{z(r)} & \dfrac{z_0^2}{z(r)^2} & -\left(\dfrac{r}{z(r)}+\sqrt{-1}\right)z_0^2 \\ -r\sqrt{-1}-z(r) & -\left(\dfrac{r}{z(r)}-\sqrt{-1}\right)z_0^2 & 1 \end{pmatrix}, \\
J &:= [\operatorname{Tr}\rho_{\theta_0} L_j L_i]_{ij} = \begin{pmatrix} 1 & -\sqrt{-1}\dfrac{z_0^2}{z(r)} \\ \sqrt{-1}\dfrac{z_0^2}{z(r)} & \dfrac{z_0^2}{z(r)^2} \end{pmatrix}, \\
\tau &:= [\operatorname{Tr}\rho_{\theta_0} L_j \sigma_i]_{ij} = \begin{pmatrix} 1 & -\sqrt{-1}\dfrac{z_0^2}{z(r)} \\ \sqrt{-1}\dfrac{z_0^2}{z(r)} & \dfrac{z_0^2}{z(r)^2} \\ -r\sqrt{-1}-z(r) & -\left(\dfrac{r}{z(r)}-\sqrt{-1}\right)z_0^2 \end{pmatrix}.
\end{aligned}$$

Given a 2×2 real positive definite matrix G, the minimal value of the weighted covariances at $\theta=\theta_0$ is given by

$$\min_{\hat{M}} \operatorname{Tr} G V_{\theta_0}[\hat{M}] = C_{\theta_0}^{(1)}\left(\rho_\theta, G\right),$$

where the minimum is taken over all estimators $\hat{M}$ that are locally unbiased at θ_0, and

$$C_{\theta_0}^{(1)}\left(\rho_\theta, G\right) = \left(\operatorname{Tr}\sqrt{\sqrt{G}J^{(S)-1}\sqrt{G}}\right)^2$$

is the Nagaoka bound [16] (see also [21]).

It can be shown that the Nagaoka bound is greater than the Holevo bound:

$$C_{\theta_0}^{(1)}\left(\rho_\theta, G\right) > C_{\theta_0}\left(\rho_\theta, G\right).$$

Let us check this fact for the special case when $G = J^{(S)}$. A direct computation shows that

$$C_{\theta_0}^{(1)}\left(\rho_\theta, J^{(S)}\right) = 4,$$

and

$$C_{\theta_0}\left(\rho_\theta, J^{(S)}\right) = \begin{cases} 2(1+z_0) - r^2(1-z_0^2), & \text{if } 0 \le r \le \sqrt{\dfrac{z_0}{1-z_0^2}} \\[2ex] 2 + \dfrac{z_0^2}{r^2(1-z_0^2)}, & \text{if } \sqrt{\dfrac{z_0}{1-z_0^2}} < r. \end{cases}$$

The right panel of Figure 1 shows the behavior of $C_{\theta_0}\left(\rho_\theta, J^{(S)}\right)$ (solid) and $C_{\theta_0}^{(1)}\left(\rho_\theta, J^{(S)}\right)$ with $z_0 = \frac{1}{4}$ (dashed) as functions of r. We see that Holevo bound $C_{\theta_0}\left(\rho_\theta, J^{(S)}\right)$ is much smaller than $C_{(0,r)}^{(1)}\left(\rho_\theta, J^{(S)}\right)$.

As in Example 16, we demonstrate that the Holevo bound is asymptotically achievable. Let

$$\Delta_i^{(n)} := \frac{1}{\sqrt{n}}\sum_{k=1}^{n} I^{\otimes k-1}\otimes L_i \otimes I^{\otimes n-k}, \qquad (i=1,2),$$

and let

$$X_j^{(n)} := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} I^{\otimes k-1} \otimes D_j \otimes I^{\otimes n-k}, \qquad (j = 1, 2, 3).$$

It then follows from the quantum central limit theorem that

$$\left(\begin{pmatrix} X^{(n)} \\ \Delta^{(n)} \end{pmatrix}, \rho_{\theta_0}^{\otimes n} \right) \underset{q}{\rightsquigarrow} N\left(0, \begin{pmatrix} \Sigma & \tau \\ \tau^* & J \end{pmatrix}\right).$$

Therefore, Corollary 11 shows that $\left(\left\{\rho_\theta^{\otimes n}\right\}, X^{(n)}\right)$ is jointly QLAN at θ_0, and that

$$(X^{(n)}, \rho_{\theta_0 + h/\sqrt{n}}^{\otimes n}) \underset{q}{\rightsquigarrow} N((\operatorname{Re}\tau)h, \Sigma)$$

for all $h \in \mathbb{R}^2$.

It should be noted that the off-diagonal block τ of the "quantum covariance" matrix is not a square matrix. This means that the derived quantum Gaussian shift model $\left\{N((\operatorname{Re}\tau)h, \Sigma)\,;\ h \in \mathbb{R}^2\right\}$ forms a submanifold of the total quantum Gaussian shift model derived in Example 16, corresponding to a 2-dimensional linear subspace in the shift parameter space. Nevertheless, Theorem 12 asserts that there exist a sequence $M^{(n)}$ of estimators on the model $\left\{\rho_{\theta_0+h/\sqrt{n}}^{\otimes n}\,;\ h \in \mathbb{R}^3\right\}$ that is asymptotically unbiased and achieves the Holevo bound:

$$\lim_{n\to\infty} \operatorname{Tr} G V_h^{(n)}[M^{(n)}] = C_h\left(N((\operatorname{Re}\tau)h, \Sigma), G\right) = C_{\theta_0}\left(\rho_\theta, G\right)$$

for all h that belong to a dense subset of $\mathbb{R}^3$.

4.3 Translating estimation of h to estimation of θ

As we have seen in the previous subsections, our theory enables us to construct asymptotically optimal estimators of h in the local models indexed by the parameter $\theta_0 + h/\sqrt{n}$. In practice of course, θ_0 is unknown and hence estimation of h, with θ_0 known, is irrelevant. The actual sequence of measurements which we have constructed depends in all interesting cases on θ_0.

However, the results immediately inspire two-step (or adaptive) procedures, in which we first measure a small proportion of the quantum systems, in number n_1 say, using some standard measurement scheme, for instance separate particle quantum tomography. From these measurement outcomes we construct an initial estimate of θ, let us call it $\widetilde{\theta}$. We can now use our theory to compute the asymptotically optimal measurement scheme which corresponds to the situation $\theta_0 = \widetilde{\theta}$. We proceed to implement this measurement on the remaining quantum systems collectively, estimating h in the model $\theta = \widetilde{\theta} + h/\sqrt{n_2}$ where n_2 is the number of systems still available for the second stage.

What can we say about such a procedure? If $n_1/n \to \alpha > 0$ as $n \to \infty$ then we can expect that the initial estimate $\widetilde{\theta}$ is root n consistent. In smooth models, one would expect that in this case the final estimate $\widehat{\theta} = \widetilde{\theta} + \widehat{h}/\sqrt{n_2}$ would be asymptotically optimal *up to a factor* $1 - \alpha$: its limiting variance will be a factor $(1-\alpha)^{-1}$ too large.

If however $n_1 \to \infty$ but $n_1/n \to \alpha = 0$ then one would expect this procedure to break down, unless the rate of growth of n_1 is very carefully chosen (and fast enough). On the other hand, instead of a direct two-step procedure, with the final estimate computed as $\widetilde{\theta} + \widehat{h}/\sqrt{n_2}$, one could be more careful in how the data obtained from the second stage measurement is used. Given the second step measurement, which results in an observed value $\widehat{h}$, one could write down the likelihood for h based on the given measurement and the initially specified model, and compute instead of the just mentioned one-step iterate, the actual maximum likelihood estimator of θ based on the second stage data. Such procedures have earlier been studied by Gill and Massar [3] and others, and shown in special cases to perform very well.

However, in general, the computational problem of even calculating the likelihood given data, measurement, and model, is challenging, due to the huge size of the Hilbert space of n copies of a finite dimensional quantum system.

5 Concluding remarks

We have developed a new theory of local asymptotic normality in a quantum regime based on a quantum extension of the log-likelihood ratio. This formulation is applicable to any model satisfying a mild smoothness condition, and is free from artificial setups such as the use of a special coordinate system and/or non-degeneracy of eigenvalues of the reference state. We also have proved asymptotic achievability of the Holevo bound for the local shift parameter on a dense subset of the parameter space.

There are of course many open questions left. Among others, it is not clear whether every sequence of statistics on a QLAN model can be realized on the limiting quantum Gaussian shift model. In classical statistics, such a problem has been solved affirmatively as the representation theorem, which asserts that, given a weakly convergent sequence $T^{(n)}$ of statistics on $\left\{p^{(n)}_{\theta_0+h/\sqrt{n}}\,;\ h\in\mathbb{R}^d\right\}$, there exist a limiting statistics T on $\left\{N(h,J^{-1})\,;\ h\in\mathbb{R}^d\right\}$ such that $T^{(n)}\overset{h}{\rightsquigarrow}T$. Representation theorem is useful in proving, for example, the non-existence of an asymptotically superefficient estimator (the converse part, as stated in Introduction). Moreover, the so-called convolution theorem and local asymptotic minimax theorem, which are the standard tools in discussing asymptotic lower bounds for estimation in LAN models, immediately follows [20]. Extending the representation theorem, convolution theorem, and local asymptotic minimax theorem to a quantum regime is an intriguing open problem. However it surely is possible to make some progress in this direction, as for instance the results of Gill and Guţă [4]. In that paper, the van Trees inequality was used to derive some results in a “poor man’s” version of QLAN theory; see also [2].

It also remains to be seen whether our asymptotically optimal statistical procedures for the local model with local parameter h can be translated into useful statistical procedures for the real world case in which θ_0 is unknown.

Appendices

A Commutation operator and the Holevo bound

In the study of quantum statistics, Holevo [9] introduced useful mathematical tools called the square summable operators and the commutation operators associated with quantum states. Let $\mathcal{H}$ be a separable Hilbert space and let ρ be a density operator. We define a real Hilbert space $\mathcal{L}^2_h(\rho)$ associated with ρ by the completion of the set $\mathcal{B}_h(\mathcal{H})$ of bounded Hermitian operators with respect to the pre-inner product $\langle X,Y\rangle_\rho := \mathrm{Re}\,\mathrm{Tr}\,\rho XY$. Letting $\rho=\sum_j s_j|\psi_j\rangle\langle\psi_j|$ be the spectral representation, an element $X\in\mathcal{L}^2_h(\rho)$ can be regarded as an equivalence class of those Hermitian operators, called the square summable operators, which satisfy $\sum_j s_j\|X\psi_j\|^2<\infty$ (so that $\psi_j\in\mathrm{Dom}(X)$ if $s_j\neq 0$) under the identification $X_1\sim X_2$ if $X_1\psi_j=X_2\psi_j$ for $s_j\neq 0$. The space $\mathcal{L}^2_h(\rho)$ thus provides a convenient tool to cope with unbounded observables. Note that when $\mathcal{H}$ is finite dimensional, the setup is considerably simplified to be $\mathcal{L}_h^2(\rho) = \mathcal{B}_h(\mathcal{H})/\ker\rho$.

Let $\mathcal{L}^2(\rho)$ be the complexification of $\mathcal{L}_h^2(\rho)$, which is also regarded as the completion of $\mathcal{B}(\mathcal{H})$ with respect to the pre-inner product

$$\langle X, Y\rangle_\rho := \frac{1}{2}\mathrm{Tr}\,\rho(YX^* + X^*Y).$$

Thus $\mathcal{L}^2(\rho)$ is a complex Hilbert space with this inner product. Let us further introduce two sesquilinear forms on $\mathcal{B}(\mathcal{H})$ by

$$(X, Y)_\rho := \mathrm{Tr}\,\rho YX^*, \qquad [X, Y]_\rho := \frac{1}{2\sqrt{-1}}\mathrm{Tr}\,\rho(YX^* - X^*Y).$$

and extend them to $\mathcal{L}^2(\rho)$ by continuity.

The *commutation operator* $\mathcal{D}_\rho : \mathcal{L}^2(\rho) \to \mathcal{L}^2(\rho)$ with respect to ρ is defined by

$$[X, Y]_\rho = \langle X, \mathcal{D}_\rho Y\rangle_\rho,$$

which is formally represented by the operator equation

$$\mathcal{D}_\rho(X)\rho + \rho\mathcal{D}_\rho(X) = \sqrt{-1}\,(X\rho - \rho X).$$

(To be precise, Holevo's original definition is different from the above one by a factor of 2.) The operator $\mathcal{D}_\rho$ is a $\mathbb{C}$-linear bounded skew-adjoint operator. Moreover, since the forms $[\,\cdot\,,\,\cdot\,]_\rho$ and $\langle\,\cdot\,,\,\cdot\,\rangle_\rho$ are real on the real subspace $\mathcal{L}_h^2(\rho)$, this subspace is invariant under the operation of $\mathcal{D}_\rho$. Thus $\mathcal{D}_\rho$ can be regarded as an $\mathbb{R}$-linear bounded skew-adjoint operator when restricted to $\mathcal{L}_h^2(\rho)$ as $\mathcal{D}_\rho : \mathcal{L}_h^2(\rho) \to \mathcal{L}_h^2(\rho)$. When no confusion is likely to arise, we drop the subscript ρ of $\mathcal{D}_\rho$ and simply denote it as $\mathcal{D}$.

Let $\mathcal{S} = \{\rho_\theta\,;\,\theta \in \Theta \in \mathbb{R}^d\}$ be a quantum statistical model satisfying the conditions: 1) the parametrization $\theta \mapsto \rho_\theta$ is smooth and nondegenerate so that the derivatives $\{\partial\rho_\theta/\partial\theta^i\}_{1\le i\le d}$ exist in trace class and form a linearly independent set at each point $\theta \in \Theta$, and 2) there exists a constant c such that

$$\left|\frac{\partial}{\partial\theta^i}\,\mathrm{Tr}\,\rho_\theta X\right|^2 \le c\,\langle X, X\rangle_{\rho_\theta}$$

for all $X \in \mathcal{B}(\mathcal{H})$ and i. The second condition assures that the linear functionals $X \mapsto (\partial/\partial\theta^i)\mathrm{Tr}\,\rho_\theta X$ can be extended to continuous linear functionals on $\mathcal{L}^2(\rho_\theta)$. Given a quantum statistical model satisfying the above conditions, the *symmetric logarithmic derivative* (SLD) $L_{\theta,i}$ in the ith direction is defined as the operator in $\mathcal{L}^2(\rho_{\rho_\theta})$ satisfying

$$\frac{\partial}{\partial\theta^i}\mathrm{Tr}\,\rho_\theta X = \langle L_{\theta,i}, X\rangle_{\rho_\theta}.$$

It is easily verified that $L_{\theta,i} \in \mathcal{L}_h^2(\rho_\theta)$; so the definition is formally written as

$$\frac{\partial\rho_\theta}{\partial\theta^i} = \frac{1}{2}(L_{\theta,i}\,\rho_\theta + \rho_\theta\,L_{\theta,i}). \tag{20}$$

When no confusion occurs, we simply denote $L_{\theta,i}$ as L_i. Since L_i is a faithful operator representation of the tangent vector $\partial/\partial\theta^i$, we shall call the $\mathbb{R}$-linear space $\mathrm{span}_{\mathbb{R}}\{L_i\}_{i=1}^d$ the *SLD tangent space* of the model ρ_θ at θ. Incidentally the $d \times d$ real symmetric matrix $J_\theta := [\mathrm{Re}\,\mathrm{Tr}\,\rho_\theta L_i L_j]_{1\le i,j\le d}$ is called the *SLD Fisher information matrix* of the model $\mathcal{S}$ at θ.

An estimator $\hat{M}$ for the parameter θ of the model $\mathcal{S}$ is called *unbiased* if

$$E_\theta[\hat{M}] = \theta \tag{21}$$

for all $\theta \in \Theta$, where $E_\theta[\,\cdot\,]$ denotes the expectation with respect to ρ_θ. An estimator $\hat{M}$ is called *locally unbiased* at $\theta_0 \in \Theta$ if the condition (21) is satisfied around $\theta = \theta_0$ up to the first order of the Taylor expansion. It is well known that an estimator $\hat{M}$ that is locally unbiased at θ_0 satisfies the quantum (SLD) Cramér-Rao inequality, $V_{\theta_0}[\hat{M}] \geq J_{\theta_0}^{-1}$, where $V_{\theta_0}[\,\cdot\,]$ denotes the covariance matrix with respect to ρ_{θ_0}. The lower bound $J_{\theta_0}^{-1}$ cannot be attained in general due to the non-commutativity of the SLDs. Because of this fact, we often switch the problem to minimizing the weighted sum of covariances, $\operatorname{Tr} GV_{\theta_0}[\hat{M}]$, given a $d \times d$ real positive definite matrix G. It is known that this quantity also has a variety of Cramér-Rao type lower bounds [9]:

$$\operatorname{Tr} GV_{\theta_0}[\hat{M}] \geq C_{\theta_0}(\rho_\theta, G)\,.$$

Among others, we concentrate our attention to the *Holevo bound* [9]:

$$\begin{aligned} C_{\theta_0}(\rho_\theta, G) \;:=\; & \min_{V,B}\{\operatorname{Tr} GV\,;\ V \text{ is a real matrix such that } V \geq Z(B),\ Z_{ij}(B) = \operatorname{Tr}\rho_{\theta_0} B_j B_i, \\ & B_1, \dots, B_d \text{ are Hermitian operators on } \mathcal{H} \text{ such that } \operatorname{Re}\operatorname{Tr}\rho_{\theta_0} L_i B_j = \delta_{ij}\}. \end{aligned} \tag{22}$$

The minimization problem over V is explicitly solved, to obtain

$$\begin{aligned} C_{\theta_0}(\rho_\theta, G) \;=\; & \min_{B}\{\operatorname{Tr} GZ(B) + \operatorname{Tr}\left|\sqrt{G}\operatorname{Im} Z(B)\sqrt{G}\right|\,;\ Z_{ij}(B) = \operatorname{Tr}\rho_{\theta_0} B_j B_i, \\ & B_1, \dots, B_d \text{ are Hermitian operators on } \mathcal{H} \text{ such that } \operatorname{Re}\operatorname{Tr}\rho_{\theta_0} L_i B_j = \delta_{ij}\}. \end{aligned}$$

Our aim here is to derive a further concise expression for it in terms of the minimal $\mathcal{D}$ invariant extension of the SLD tangent space.

Theorem 19. *Given a quantum statistical model $\{\rho_\theta\,;\ \theta \in \Theta \subset \mathbb{R}^d\}$ on $\mathcal{H}$, let $\mathcal{T}$ be the minimal $\mathcal{D}$ invariant extension of the SLD tangent space $\operatorname{span}_{\mathbb{R}}\{L_i\}_{i=1}^d$ of the model at $\theta = \theta_0$, and let $\{D_j\}_{j=1}^r$ be a basis of $\mathcal{T}$. The Holevo bound defined by (22) is rewritten as*

$$\begin{aligned} C_{\theta_0}(\rho_\theta, G) \;=\; & \min_{F}\{\operatorname{Tr} GZ + \operatorname{Tr}\left|\sqrt{G}\operatorname{Im} Z\sqrt{G}\right|\,;\ Z = {}^t F \Sigma F, \\ & F \text{ is an } r \times d \text{ real matrix satisfying } {}^t F \operatorname{Re}(\tau) = I\}, \end{aligned} \tag{23}$$

where Σ and τ are $r \times r$ and $r \times d$ complex matrices whose (i,j)th entries are given by $\Sigma_{ij} = \operatorname{Tr}\rho_{\theta_0} D_j D_i$ and $\tau_{ij} = \operatorname{Tr}\rho_{\theta_0} L_j D_i$.

Proof. Let $\mathcal{T}^\perp$ be the orthogonal complement of $\mathcal{T}$ in $\mathcal{L}_h^2(\rho_{\theta_0})$ with respect to the inner product $\langle\cdot,\cdot\rangle_{\rho_{\theta_0}}$, and let $\mathcal{P} : \mathcal{L}_h^2(\rho_{\theta_0}) \to \mathcal{T}$ and $\mathcal{P}^\perp : \mathcal{L}_h^2(\rho_{\theta_0}) \to \mathcal{T}^\perp$ be the projections associated with the decomposition $\mathcal{L}_h^2(\rho_{\theta_0}) = \mathcal{T} \oplus \mathcal{T}^\perp$. Note that if $X \in \mathcal{T}^\perp$ and $Y \in \mathcal{T}$, then

$$(X, Y)_{\rho_{\theta_0}} = \langle X, Y\rangle_{\rho_{\theta_0}} + \sqrt{-1}\,\langle X, \mathcal{D}Y\rangle_{\rho_{\theta_0}} = 0.$$

We show that the operators $\{B_j\}_{j=1}^d$ that achieve the minimum in (22) can be taken from $\mathcal{T}$. Let $\{B_j\}_{j=1}^d \subset \mathcal{L}_h^2(\rho_{\theta_0})$ satisfies the local unbiasedness condition $\operatorname{Re}\operatorname{Tr}\rho_{\theta_0} L_i B_j = \delta_{ij}$, which is rewritten as

$$\langle L_i, B_j\rangle_{\rho_{\theta_0}} = \delta_{ij}.$$

Then $\{\mathcal{P}(B_j)\}_{j=1}^d$ also satisfies the local unbiasedness

$$\langle L_i, \mathcal{P}(B_j)\rangle_{\rho_{\theta_0}} = \langle L_i, B_j\rangle_{\rho_{\theta_0}} = \delta_{ij}.$$

Further,

$$\begin{aligned} Z_{ij}(B) &= (B_i, B_j)_{\rho_{\theta_0}} = \left(\mathcal{P}(B_i) + \mathcal{P}^{\perp}(B_i), \mathcal{P}(B_j) + \mathcal{P}^{\perp}(B_j)\right)_{\rho_{\theta_0}} \\ &= (\mathcal{P}(B_i), \mathcal{P}(B_j))_{\rho_{\theta_0}} + \left(\mathcal{P}^{\perp}(B_i), \mathcal{P}^{\perp}(B_j)\right)_{\rho_{\theta_0}} = Z_{ij}(\mathcal{P}(B)) + Z_{ij}(\mathcal{P}^{\perp}(B)). \end{aligned}$$

Since $Z(\,\cdot\,)$ is a Gram matrix and is positive semidefinite, this decomposition implies that $Z(B) \geq Z(\mathcal{P}(B))$. Thus the observables B that minimize (22) can be taken from $\mathcal{T}$.

Let $B_j \in \mathcal{T}$ be expanded as $B_j = F_j^k D_k$, where F is an $r \times d$ real matrix. Then the local unbiasedness condition is rewritten as

$$\langle L_i, B_j \rangle_{\rho_{\theta_0}} = F_j^k \, \langle L_i, D_k \rangle_{\rho_{\theta_0}} = \delta_{ij},$$

or in a matrix form,

$${}^t F \, (\operatorname{Re} \tau) = I.$$

Further, the Gram matrix $Z(B)$ is rewritten as

$$Z_{ij}(B) = (B_i, B_j)_{\rho_{\theta_0}} = F_i^k F_j^\ell \, (D_k, D_\ell)_{\rho_{\theta_0}} \,,$$

or,

$$Z(B) = {}^t F \Sigma F.$$

This proves the claim. $\square$

When the SLD tangent space itself is $\mathcal{D}$ invariant, the Holevo bound can be represented in terms of the RLD Fisher information matrix as follows.

Corollary 20. *Let $\left\{\rho_\theta \,;\, \theta \in \Theta \subset \mathbb{R}^d\right\}$ be a quantum statistical model, and let L_i $(1 \leq i \leq d)$ be the SLDs at θ_0. If the SLD tangent space $\operatorname{span}_{\mathbb{R}} \{L_i\}_{i=1}^d$ at θ_0 is $\mathcal{D}$ invariant, then*

$$C_{\theta_0}(\rho_\theta, G) = \operatorname{Tr} G (J^{(R)})^{-1} + \operatorname{Tr} \left| \sqrt{G} \operatorname{Im} (J^{(R)})^{-1} \sqrt{G} \right|,$$

where $(J^{(R)})^{-1} := (\operatorname{Re} J)^{-1} J \, (\operatorname{Re} J)^{-1}$ with $J_{ij} = \operatorname{Tr} \rho_{\theta_0} L_j L_i$.

Proof. Let us set $D_i := L_i$ for $1 \leq i \leq d$ in Theorem 19. Then $\Sigma = \tau$, and the local unbiasedness condition ${}^t F \, (\operatorname{Re} \tau) = I$ has a unique solution $F = (\operatorname{Re} \Sigma)^{-1}$, whereby $Z = (\operatorname{Re} J)^{-1} J \, (\operatorname{Re} J)^{-1}$. $\square$

Note that RLDs may not exist if the model is degenerate (i.e., non-faithful). This means that $J^{(R)}$ may not be well-defined for such a model. Nevertheless we use the notation $(J^{(R)})^{-1}$ even for a degenerate model, and call it the inverse of the RLD Fisher information matrix, as long as the SLD tangent space is $\mathcal{D}$ invariant. For an idea behind this nomenclature, consult [1].

Finally, we show that the Holevo bound for the nth i.i.d. extension model is precisely $\frac{1}{n}$ times that for the base model.

Corollary 21. *Given a quantum statistical model $\mathcal{S} = \left\{\rho_\theta \,;\, \theta \in \Theta \subset \mathbb{R}^d\right\}$, let $\mathcal{S}^{(n)} = \left\{\rho_\theta^{\otimes n} \,;\, \theta \in \Theta \subset \mathbb{R}^d\right\}$ be the nth i.i.d. extension model. Then*

$$C_{\theta_0}\left(\rho_\theta^{\otimes n}, G\right) = \frac{1}{n} C_{\theta_0}(\rho_\theta, G) \,.$$

Proof. Let us distinguish quantities that belong to models of different extension by specifying the degree n of extension in the superscript. Letting $\{L_i\}_{i=1}^d$ and $\{D_j\}_{j=1}^r$ be SLDs and a basis of $\mathcal{T}$ in Theorem 19, the corresponding quantities for $\mathcal{S}^{(n)}$ are given by

$$L_i^{(n)} = \sum_{k=1}^{n} I^{\otimes k-1} \otimes L_i \otimes I^{\otimes n-k}$$

and

$$D_j^{(n)} = \sum_{k=1}^{n} I^{\otimes k-1} \otimes D_j \otimes I^{\otimes n-k}.$$

Thus

$$\Sigma^{(n)} = n\Sigma^{(1)}, \quad \tau^{(n)} = n\tau^{(1)}, \quad F^{(n)} = \frac{1}{n}F^{(1)},$$

so that

$$Z^{(n)} = {}^tF^{(n)}\Sigma^{(n)}F^{(n)} = \frac{1}{n}Z^{(1)},$$

and

$$C_{\theta_0}\left(\rho_\theta^{\otimes n}, G\right) = \frac{1}{n}\,C_{\theta_0}\left(\rho_\theta, G\right)$$

doe to Theorem 19. □

B Estimation of quantum Gaussian shift model

In this section, we briefly overview the estimation theory for a quantum Gaussian shift model. For a mathematically rigorous treatment, consult [9].

Lemma 22. *Let* $(X, \phi_h) \sim N(h, J)$*, where* J *is a* $d \times d$ *positive semidefinite complex matrix. Then*

$$\phi_h(X_i) = h_i \tag{24}$$

and

$$\phi_h((X_j - h_j)(X_i - h_i)) = J_{ij} \tag{25}$$

hold.

Proof. Letting $U(\xi) := \mathrm{e}^{\sqrt{-1}\xi^i X_i}$,

$$\begin{aligned} \phi_h(U(\xi)) &= 1 + \sqrt{-1}\phi_h(\xi^i X_i) - \frac{1}{2}\phi_h(\left(\xi^i X_i\right)^2) + o(\xi^2) \\ &= 1 + \sqrt{-1}\phi_h(X_i)\xi^i - \frac{1}{2}\phi_h(X_i X_j)\xi^i\xi^j + o(\xi^2) \\ &= 1 + \sqrt{-1}\phi_h(X_i)\xi^i - \frac{1}{2}\phi_h(X_i \circ X_j)\xi^i\xi^j + o(\xi^2), \end{aligned}$$

where $X_i \circ X_j = \frac{1}{2}\left(X_i X_j + X_J X_i\right)$. Further, letting $V = \operatorname{Re} J$ and $S = \operatorname{Im} J$,

$$\begin{aligned} \mathrm{e}^{\sqrt{-1}\xi^i h_i - \frac{1}{2}V_{ij}\xi^i\xi^j} &= 1 + \left(\sqrt{-1}\xi^i h_i - \frac{1}{2}V_{ij}\xi^i\xi^j\right) + \frac{1}{2}\left(\sqrt{-1}\xi^i h_i - \frac{1}{2}V_{ij}\xi^i\xi^j\right)^2 + o(\xi^2) \\ &= 1 + \sqrt{-1}\xi^i h_i - \frac{1}{2}\left(V_{ij} + h_i h_j\right)\xi^i\xi^j + o(\xi^2). \end{aligned}$$

A comparison immediately leads to (24) and the identity $\phi_h(X_i \circ X_j) = V_{ij} + h_i h_j$. Thus

$$\begin{aligned} \phi_h((X_j - h_j)(X_i - h_i)) &= \phi_h(X_j X_i - h_j X_i - h_i X_j + h_i h_j) \\ &= \phi_h(X_j X_i) - h_i h_j \\ &= \phi_h\left(X_i \circ X_j - \frac{1}{2}[X_i, X_j]\right) - h_i h_j = J_{ij}. \end{aligned}$$

□

In what follows, we treat the quantum Gaussian shift model $\left\{N(\tau h, \Sigma);\ h \in \mathbb{R}^d\right\}$ on $\mathrm{CCR}\,(\mathrm{Im}\,\Sigma)$, where Σ is an $r \times r$ complex matrix such that $\Sigma \geq 0$ and $\mathrm{Re}\,\Sigma > 0$, and τ is an $r \times d$ real matrix with $d \leq r$ such that $\mathrm{rank}\,\tau = d$. Let $X = (X_1, \ldots, X_r)$ be the basic canonical observables of $\mathrm{CCR}\,(\mathrm{Im}\,\Sigma)$, and $(X, \phi_h) \sim N(\tau h, \Sigma)$.

Lemma 23. *Let $U(\xi) := \mathrm{e}^{\sqrt{-1}\xi^i X_i}$. The SLD L_i $(1 \leq i \leq d)$ at h defined by*

$$\frac{\partial}{\partial h_k}\phi_h(U(\xi)) = \frac{1}{2}\phi_h\left(U(\xi)L_k + L_k U(\xi)\right) \tag{26}$$

is given by

$$L_k = \sum_{\ell=1}^{r}\left[(\mathrm{Re}\,\Sigma)^{-1}\tau\right]_{\ell k}(X_\ell - (\tau h)_\ell I). \tag{27}$$

Proof. In this proof we lift Einstein's summation convention. Let $V = \mathrm{Re}\,\Sigma$ and $S = \mathrm{Im}\,\Sigma$, and fix a $k \in \{1, \ldots, d\}$ arbitrarily. Due to the Baker-Hausdorff formula,

$$U(\xi) = \mathrm{e}^{\sqrt{-1}\sum_{i=1}^r \xi^i X_i} = \exp\left(-\sqrt{-1}\sum_{i=1}^{r} S_{ki}\xi^k\xi^i\right)\exp\left(\sqrt{-1}\xi^k X_k\right)\exp\left(\sqrt{-1}\sum_{i\neq k}\xi^i X_i\right).$$

By differentiating in ξ^k, we have

$$\frac{\partial}{\partial \xi^k}U(\xi) = -\sqrt{-1}\left(\sum_{i=1}^{r} S_{ki}\xi^i - X_k\right)U(\xi).$$

Thus

$$\begin{aligned} \phi_h((X_k - (\tau h)_k I)U(\xi)) &= \phi_h\left(\left(\sum_{i=1}^{r} S_{ki}\xi^i - \sqrt{-1}\frac{\partial}{\partial \xi^k} - (\tau h)_k I\right)U(\xi)\right) \\ &= \left(\sum_{i=1}^{r} S_{ki}\xi^i - \sqrt{-1}\frac{\partial}{\partial \xi^k} - (\tau h)_k\right)\phi_h(U(\xi)) \\ &= \left(\sum_{i=1}^{r} S_{ki}\xi^i - \sqrt{-1}\frac{\partial}{\partial \xi^k} - (\tau h)_k\right)\mathrm{e}^{\sqrt{-1}\,{}^t\xi\tau h - \frac{1}{2}\,{}^t\xi V\xi} \\ &= \left(\sum_{i=1}^{r} S_{ki}\xi^i - (\tau h)_k\right)\phi_h(U(\xi)) - \sqrt{-1}\left(\sqrt{-1}(\tau h)_k - (V\xi)_k\right)\phi_h(U(\xi)) \\ &= \left(S\xi + \sqrt{-1}V\xi\right)_k \phi_h(U(\xi)) \\ &= \sqrt{-1}\left(\bar{J}\xi\right)_k \phi_h(U(\xi)). \end{aligned} \tag{28}$$

Similarly, we obtain

$$\phi_h(U(\xi)(X_k - (\tau h)_k I)) = \sqrt{-1}\left(J\xi\right)_k \phi_h(U(\xi)). \tag{29}$$

By combining (28) and (29),

$$\phi_h\left((X_k - (\tau h)_k I)U(\xi) + U(\xi)(X_k - (\tau h)_k I) \right) = 2\sqrt{-1}\,(V\xi)_k\,\phi_h(U(\xi)). \tag{30}$$

On the other hand, by a direct calculation

$$\frac{\partial}{\partial h_k}\phi_h(U(\xi)) = \frac{\partial}{\partial h_k}\mathrm{e}^{\sqrt{-1}\,{}^t\xi\tau h - \frac{1}{2}\,{}^t\xi V\xi} = \sqrt{-1}({}^t\xi\tau)_k\phi_h(U(\xi)). \tag{31}$$

A comparison between (30) and (31) yields

$$L_k = \sum_{\ell=1}^{r}\left[V^{-1}\tau\right]_{\ell k}(X_\ell - (\tau h)_\ell I).$$

$\square$

Let $\tilde{L}_k := X_k - (\tau h)_k I$. It follows from (28) and (29) that $\mathcal{D}_{\phi_h}(\tilde{L}_i) = \sum_{i=1}^{r}(V^{-1}S)_{ki}\tilde{L}_k$, where $\mathcal{D}_{\phi_h}$ is the commutation operator with respect to ϕ_h defined by

$$\phi_h\left(U(\xi)\mathcal{D}_{\phi_h}(X) + \mathcal{D}_{\phi_h}(X)U(\xi)\right) = \sqrt{-1}\phi_h\left(U(\xi)X - XU(\xi)\right).$$

This means $\mathcal{T} = \operatorname{span}\left\{\tilde{L}_k\right\}_{k=1}^{r}$ is $\mathcal{D}_{\phi_h}$ invariant. Further, we can check from (27) that $\operatorname{span}\{L_i\}_{i=1}^{d} \subset \mathcal{T}$ and

$$\phi_h(\tilde{L}_j\tilde{L}_i) = \Sigma_{ij} \tag{32}$$

and

$$\operatorname{Re}\phi_h(L_j\tilde{L}_i) = \tau_{ij}. \tag{33}$$

These relations play a fundamental role in connecting a general quantum statistical model $\mathcal{S} = \left\{\rho_\theta\,;\,\theta \in \Theta \subset \mathbb{R}^d\right\}$ on $\mathcal{H}$ with a quantum Gaussian shift model $\mathcal{G} = \left\{N(\tau h, \Sigma)\,;\,h \in \mathbb{R}^d\right\}$ as follows. Let $\{L_i^{\mathcal{S}}\}_{i=1}^{d}$ be the SLDs of the model $\mathcal{S}$ at $\theta = \theta_0$, and let $\mathcal{T}^{\mathcal{S}}$ the minimal $\mathcal{D}^{\mathcal{S}}$ invariant extension of the SLD tangent space $\operatorname{span}\{L_i^{\mathcal{S}}\}_{i=1}^{d}$. Further let $\{D_j^{\mathcal{S}}\}_{j=1}^{r}$ be a basis of $\mathcal{T}^{\mathcal{S}}$ and let Σ and τ are $r \times r$ and $r \times d$ matrices whose (i,j)th entries are given by $\Sigma_{ij} = \operatorname{Tr}\rho_{\theta_0}D_jD_i$ and $\tau_{ij} = \operatorname{Re}\operatorname{Tr}\rho_{\theta_0}L_jD_i$. Based on those information, we introduce a quantum Gaussian shift model $\mathcal{G} = \left\{N(\tau h, \Sigma)\,;\,h \in \mathbb{R}^d\right\}$ on $\mathrm{CCR}\,(\operatorname{Im}\Sigma)$, which exhibits relations (32) and (33). Recall that the Holevo bound of a quantum statistical model is completely determined by the information Σ and τ (Theorem 19). We thus obtain the following important consequence.

Corollary 24. *The Holevo bound $C_{\theta_0}(\rho_\theta, G)$ for the model $\mathcal{S}$ at $\theta = \theta_0$ is identical to the Holevo bound $C_h(N(\tau h, \Sigma), G)$ for the Gaussian shift model $\mathcal{G}$.*

As to the achievability of the Holevo bound $C_h(N(\tau h, \Sigma), G)$ for the Gaussian shift model $\mathcal{G}$, we have the following.

Theorem 25. *Given a weight $G > 0$, there exist an unbiased estimator $\hat{M}$ that achieves the Holevo bound for the model $\{N(\tau h, \Sigma)\,;\,h \in \mathbb{R}^d\}$, i.e.,*

$$\operatorname{Tr} GV_h[\hat{M}] = C_h\left(N(\tau h, \Sigma), G\right).$$

Proof. Let F be the matrix that achieve the minimum of (23) for the model $\{N(\tau h, \Sigma)\}_h$, and let $Z = {}^tF\Sigma F$. Further, let $\tilde{V} = \operatorname{Re} Z$, $\tilde{S} = \operatorname{Im} Z$. $\hat{V} = \sqrt{G^{-1}}\left|\sqrt{G}\operatorname{Im} Z\sqrt{G}\right|\sqrt{G^{-1}}$, and $\hat{Z} = \hat{V} - \sqrt{-1}\tilde{S}$.

We introduce an ancillary quantum Gaussian state $(Y, \psi) \sim N(0, \hat{Z})$ on another CCR $\left(-\tilde{S}\right)$, and a set of canonical observables

$$\overline{X}_i := \tilde{X}_i \otimes I + I \otimes Y_i \quad (1 \le i \le d),$$

on CCR $\left(\tilde{S}\right) \otimes$ CCR $\left(-\tilde{S}\right)$, where $\tilde{X}_i = F_i^k X_k$. It is important to notice that the CCR subalgebra $\mathcal{A}[\overline{X}]$ generated by $\{\overline{X}_i\}_{1\le i \le d}$ is a commutative one because

$$\frac{\sqrt{-1}}{2}[\overline{X}_i, \overline{X}_j] = \tilde{S}_{ij} - \tilde{S}_{ij} = 0$$

for $1 \le i, j \le d$. Moreover

$$(\phi_h \otimes \psi)(\mathrm{e}^{\sqrt{-1}\xi^i \overline{X}_i}) = \left[\phi_h\left(\mathrm{e}^{\sqrt{-1}\xi^i \tilde{X}_i}\right)\right]\left[\psi\left(\mathrm{e}^{\sqrt{-1}\xi^i Y_i}\right)\right] = \mathrm{e}^{\sqrt{-1}\xi^i h_i - \frac{1}{2}\xi^i\xi^j(\tilde{V}+\hat{V})_{ij}}.$$

This means that the observables $\overline{X}_i$ $(1 \le i \le d)$ follow the classical Gaussian distribution $N(h, \tilde{V} + \hat{V})$. In particular,

$$E_h[\overline{X}] = h$$

for all $h \in \mathbb{R}^d$, and

$$\operatorname{Tr} GV_h[\overline{X}] = \operatorname{Tr} G(\tilde{V} + \hat{V}) = C_h\left(N(\tau h, \Sigma), G\right).$$

The claim was verified. $\square$

C Estimation theory for pure state models

Lemma 26. *Let ρ be a pure state and $A_1, \ldots, A_d$ observables on a finite dimensional Hilbert space $\mathcal{H}$. If $J_{ij} := \operatorname{Tr} \rho A_j A_i$ are all real for $1 \le i, j \le d$, there exist observables $K_1, \ldots, K_d$ such that*

$$[A_i + K_i, A_j + K_j] = 0,$$

for $1 \le i, j \le d$ and

$$K_i \rho = 0$$

for $1 \le i \le d$.

Proof. Let $\rho := |\psi\rangle\langle\psi|$, and let $|l_i\rangle := A_i|\psi\rangle$ for $1 \le i \le d$. Because $\langle\psi|l_i\rangle$ and $\langle l_i|l_j\rangle$ $(= J_{ji})$ are all real, there exist a CONS $\{|e_k\rangle\}_{k=1}^{\dim\mathcal{H}}$ of $\mathcal{H}$ such that $\langle e_k|\psi\rangle$ and $\langle e_k|l_i\rangle$ are all real, and that $\langle e_k|\psi\rangle \ne 0$ for all k. Let

$$\tilde{A}_i := \sum_{k=1}^{\dim\mathcal{H}} \frac{\langle e_k|l_i\rangle}{\langle e_k|\psi\rangle}|e_k\rangle\langle e_k|,$$

and $K_i := \tilde{A}_i - A_i$. Obviously $[A_i + K_i, A_j + K_j] = [\tilde{A}_i, \tilde{A}_j] = 0$, and

$$K_i|\psi\rangle = \left(\tilde{A}_i - A_i\right)|\psi\rangle = |l_i\rangle - |l_i\rangle = 0.$$

This means $K_i\rho = 0$. $\square$

Theorem 27. *Let $\left\{\rho_\theta\,;\ \theta \in \Theta \subset \mathbb{R}^d\right\}$ be a quantum statistical model comprising pure states on a finite dimensional Hilbert space $\mathcal{H}$, and let $C_{\theta_0}\left(\rho_\theta, G\right)$ be the Holevo bound at $\theta_0 \in \Theta$ for a given weight $G > 0$. There exist a locally unbiased estimator $\hat{M}$ at $\theta_0 \in \Theta$ such that $\operatorname{Tr} GV[\hat{M}] = C_{\theta_0}\left(\rho_\theta, G\right)$.*

Proof. Let $\mathcal{T}$ be the minimal $\mathcal{D}$ invariant extension of the SLD tangent space $\operatorname{span}\{L_i\}_{i=1}^{d}$ of the model $\{\rho_\theta\}$ at $\theta = \theta_0$, i.e., containing all the SLDs $\{L_i\}_{i=1}^{d}$ of $\{\rho_\theta\}$ at θ_0, let $\{D_j\}_{j=1}^{r}$ be a basis of $\mathcal{T}$. Let Σ, τ be $r \times r$, $r \times d$ complex matrices defined by $\Sigma_{ij} = \operatorname{Tr} \rho_{\theta_0} D_j D_i$, $\tau_{ij} = \operatorname{Tr} \rho_{\theta_0} L_j D_i$. According to Theorem 19, the Holevo bound for a weight $G > 0$ can be expressed

$$\begin{aligned} C_{\theta_0}(\rho_\theta, G) \quad &= \quad \min_F \{\operatorname{Tr} GZ + \operatorname{Tr} \left|\sqrt{G} \operatorname{Im} Z \sqrt{G}\right| \; ; \; Z = {}^t F \Sigma F, \\ &\qquad F \text{ is an } r \times d \text{ real matrix satisfying } {}^t F \operatorname{Re}(\tau) = I\}. \end{aligned} \tag{34}$$

Let F be the matrix that attains the minimum in (34), and let $Z := {}^t F \Sigma F$, $\tilde{V} := \operatorname{Re} Z$, $\tilde{S} := \operatorname{Im} Z$, $\hat{V} = \sqrt{G^{-1}} \left|\sqrt{G} \operatorname{Im} Z \sqrt{G}\right| \sqrt{G^{-1}}$, and $\hat{Z} = \hat{V} - \sqrt{-1}\tilde{S}$. Lemma 13 assures that there exist a Hilbert space $\hat{\mathcal{H}}$ and a pure state σ and observables B_i $(1 \le i \le d)$ on $\hat{\mathcal{H}}$ such that $\operatorname{Tr} \sigma B_i = 0$ and $\operatorname{Tr} \sigma B_j B_i = \hat{Z}_{ij}$. Further, let

$$\overline{X}_i := \tilde{X}_i \otimes \hat{I} + I \otimes B_i \quad (1 \le i \le d),$$

where $\tilde{X}_i := F_i^k D_k$ $(1 \le i \le d)$, and $\hat{I}$ is the identity matrix on $\hat{\mathcal{H}}$. It then follows that

$$\operatorname{Tr}\,(\rho_{\theta_0} \otimes \sigma)\, \overline{X}_j \overline{X}_i = \left(\tilde{V} + \hat{V}\right)_{ij}. \tag{35}$$

According to Lemma 26, there exist observables $K_1, \ldots, K_d$ on $\mathcal{H} \otimes \hat{\mathcal{H}}$ such that $[\overline{X}_i + K_i, \overline{X}_j + K_j] = 0$ and $K_i\,(\rho_{\theta_0} \otimes \sigma) = 0$. Let $\hat{T}_i := \theta_0^i I \otimes \hat{I} + \left(\overline{X}_i + K_i\right)$. Then $\hat{T}_1, \ldots, \hat{T}_d$ are simultaneously measurable, and satisfy the local unbiasedness condition:

$$\operatorname{Tr}\,(\rho_{\theta_0} \otimes \sigma)\, \hat{T}_j = \theta_0^j$$

and

$$\begin{aligned} \operatorname{Tr}\,(\partial_i \rho_{\theta_0} \otimes \sigma)\, \hat{T}_j \quad &= \quad \operatorname{Tr} \partial_i \rho_{\theta_0} \tilde{X}_j \\ &= \quad F_j^k \operatorname{Tr} \partial_i \rho_{\theta_0} D_k \\ &= \quad F_j^k \operatorname{Re} \operatorname{Tr} \rho_{\theta_0} L_i D_k \\ &= \quad \{F\,(\operatorname{Re} \tau)\}_{ji} = \delta_{ij}. \end{aligned}$$

Further

$$V_{\theta_0}[\hat{T}]_{ij} = \operatorname{Tr}\,(\rho_{\theta_0} \otimes \sigma) \left(\overline{X}_i + K_i\right) \left(\overline{X}_i + K_i\right) = \left(\tilde{V} + \hat{V}\right)_{ij}.$$

This completes the proof. □

D Quantum central limit theorem

Jakšić, Pautrat, and Pillet [10] proved the following strong version of a quantum central limit theorem.

Proposition 28. *Given a sequence $\mathcal{H}^{(n)}$ of Hilbert space, let $\rho^{(n)}$ and $A^{(n)} = (A_1^{(n)}, \ldots, A_d^{(n)})$ be a state and a list of observables on $\mathcal{H}^{(n)}$ that enjoy the quantum central limit theorem in the sense of convergence of the quasi-characteristic function:*

$$\left(A^{(n)}, \rho^{(n)}\right) \underset{q}{\rightsquigarrow} N(h, J) \sim (X, \phi),$$

where J is a $d \times d$ positive semidefinite matrix. Then for any bounded continuous functions $f_1, \ldots, f_m$ and a noncommutative polynomial P, it follows that

$$\lim_{n \to \infty} \operatorname{Tr} \rho^{(n)} P\left(\overrightarrow{f(A^{(n)})}\right) = \phi\left(P\left(\overrightarrow{f(X)}\right)\right),$$

where $\overrightarrow{f(B)} := (f_1(B_1), \ldots, f_1(B_d), \ldots, f_m(B_1), \ldots, f_m(B_d))$ *for a given list* $B = (B_1, \ldots, B_d)$ *of observables, and* $P\left(\overrightarrow{f(B)}\right) := P(f_1(B_1), \ldots, f_1(B_d), \ldots, f_m(B_1), \ldots, f_m(B_d))$.

Proposition 28 is strong enough to prove the following, which is essential in constructing a sequence of POVMs that asymptotically achieves the Holevo bound (Theorem 12).

Corollary 29. *Under the same assumption as in Proposition 28, for any bounded continuous functions* $g, f_1, \ldots, f_m$*, and noncommutative polynomials* P, Q*, with* P *being Hermitian operator-valued, it follows that*

$$\begin{aligned}\lim_{n\to\infty} \operatorname{Tr} \rho^{(n)} g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) Q\left(\overrightarrow{f(A^{(n)})}\right) g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) \\ = \phi\left(g\left(P\left(\overrightarrow{f(X)}\right)\right) Q\left(\overrightarrow{f(X)}\right) g\left(P\left(\overrightarrow{f(X)}\right)\right)\right).\end{aligned}$$

Proof. Let $l := \max_{1\le i\le m} \sup_x |f_i(x)|$. There exist $l_P > 0$ and $l_Q > 0$ such that $l_P > \left\|P(\overrightarrow{B})\right\|$ and $l_Q > \left\|Q(\overrightarrow{B})\right\|$ for any list $\overrightarrow{B} = (B_1, \ldots, B_{dm})$ of observables such that $\|B_i\| \le l$. Let $l_g := \sup\{|g(x)|\ ;\ x \in [-l_P, l_P]\}$. There exist a sequence $R^{(k)}(x)$ of polynomials that uniformly converges to $g(x)$ on $[-l_P, l_P]$.

Let

$$a_{kn} := \operatorname{Tr} \rho^{(n)} R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) Q\left(\overrightarrow{f(A^{(n)})}\right) R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right),$$

and let

$$a_n := \operatorname{Tr} \rho^{(n)} g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) Q\left(\overrightarrow{f(A^{(n)})}\right) g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right).$$

We show that a_{kn} uniformly converges to a_n as $k \to \infty$. In fact, letting $l_R := \sup\left\{R^{(k)}(x)\ ;\ k \in \mathbb{N}, x \in [-l_P, l_P]\right\}$,

$$\begin{aligned}
&\sup_{n\in\mathbb{N}} |a_n - a_{kn}| \\
&\quad = \sup_{n\in\mathbb{N}} \Bigg| \operatorname{Tr} \rho^{(n)} g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) Q\left(\overrightarrow{f(A^{(n)})}\right) g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) \\
&\quad = -\operatorname{Tr} \rho^{(n)} R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) Q\left(\overrightarrow{f(A^{(n)})}\right) R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) \Bigg| \\
&\quad \le \sup_{n\in\mathbb{N}} \left| \operatorname{Tr} \rho^{(n)} g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) Q\left(\overrightarrow{f(A^{(n)})}\right) \left[g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) - R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right)\right] \right| \\
&\qquad + \sup_{n\in\mathbb{N}} \left| \operatorname{Tr} \rho^{(n)} \left[g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) - R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right)\right] Q\left(\overrightarrow{f(A^{(n)})}\right) R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) \right| \\
&\quad \le l_g l_Q \sup_{n\in\mathbb{N}} \left\| g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) - R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) \right\| \\
&\qquad + l_Q l_R \sup_{n\in\mathbb{N}} \left\| g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) - R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) \right\| \\
&\quad \le l_Q(l_g + l_R) \sup_{x\in[-l_P, l_P]} \left| g(x) - R^{(k)}(x) \right|,
\end{aligned}$$

which converges to zero as $k \to \infty$.

The uniform convergence $a_{kn} \rightrightarrows a_n$ as well as the existence of $\lim_{k\to\infty} \lim_{n\to\infty} a_{kn}$, which follows from Proposition 28, ensure that

$$
\begin{aligned}
&\lim_{n\to\infty} \operatorname{Tr} \rho^{(n)} g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) Q\left(\overrightarrow{f(A^{(n)})}\right) g\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) \\
&\quad = \lim_{n\to\infty} \lim_{k\to\infty} \operatorname{Tr} \rho^{(n)} R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) Q\left(\overrightarrow{f(A^{(n)})}\right) R^{(k)}\left(P\left(\overrightarrow{f(A^{(n)})}\right)\right) \\
&\quad = \lim_{n\to\infty} \lim_{k\to\infty} a_{kn} \\
&\quad = \lim_{k\to\infty} \lim_{n\to\infty} a_{kn} \\
&\quad = \lim_{k\to\infty} \phi\left(R^{(k)}\left(P\left(\overrightarrow{f(X)}\right)\right) Q\left(\overrightarrow{f(X)}\right) R^{(k)}\left(P\left(\overrightarrow{f(X)}\right)\right)\right) \\
&\quad = \phi\left(g\left(P\left(\overrightarrow{f(X)}\right)\right) Q\left(\overrightarrow{f(X)}\right) g\left(P\left(\overrightarrow{f(X)}\right)\right)\right).
\end{aligned}
$$

This proves the claim. □

References

[1] Fujiwara, A. and Nagaoka, H. (1999). An estimation theoretical characterization of coherent states. *J. Math. Phys.* **40**, 4227.

[2] Gill, R. D. and Levit, B. Y. (1995). Applications of the Van Trees inequality: A Bayesian Cramér-Rao bound. *Bernoulli* **1**, 59-79.

[3] Gill, R. D. and Massar, S. (2000). State estimation for large ensembles. *Physical Review A*, **61**, 042312.

[4] Gill, R. D. and Guţă, M. (2012). On Asymptotic Quantum Statistical Inference. *IMS Collections From Probability to Statistics and Back: High-Dimensional Models and Processes* Vol. 9 105-127.

[5] Guţă, M. and Kahn, J. (2006). Local asymptotic normality for qubit states. *Phys. Rev. A*, 73:052108.

[6] Guţă, M and Jençová, A. (2007). Local asymptotic normality in quantum statistics. *Commun. Math. Phys., 276*, 341–379.

[7] Hayashi, M. (1997). A linear programming approach to attainable Cramér-Rao type bounds. in *Quantum Communication, Computing, and Measurement*, ed. by Hirota et al. Plenum, NY, 99-108.

[8] Hayashi, M. and Matsumoto, K. (2008). Asymptotic performance of optimal state estimation in qubit system. *J. Math. Phys.* **49**, 102101.

[9] Holevo, A. S. (1982). *Probabilistic and Statistical Aspects of Quantum Theory.* North-Holland, Amsterdam. First appeared, in Russian, 1980.

[10] Jakšić, V., Pautrat, Y., and Pillet, C.-A. (2010). A quantum central limit theorem for sums of independent identically distributed random variables. *J. Math. Phys.* **51**, 015208.

[11] Kahn, J. and Guţă, M. (2009) Local asymptotic normality for finite dimensional quantum systems. *Commun. Math. Phys. 289*, 597-652.

[12] Kuperberg, G. (2005). A tracial quantum central limit theorem. *Trans. Am. Math. Soc.* **357**, 459.

[13] Le Cam, L. (1986). *Asymptotic Methods in Statistical Decision Theory*, Springer Verlag, New York.

[14] Manuceau, J., Sirugue, M., Testard, D., and Verbeure, A. (1973). The smallest C-algebra for canonical commutations relations. *Commun. Math. Phys.* **32**, 231.

[15] Matsumoto, K. (2002). A new approach to the Cramer-Rao type bound of the pure state model. *J. Phys. A* **35**, 3111.

[16] Nagaoka, H. (1991). A generalization of the simultaneous diagonalization of Hermitian matrices and its relation to quantum estimation theory. *Transactions of the Japan Society for Industrial and Applied Mathematics*, **1**, 305-318 (in Japanese).

[17] Ohya, M. and Petz, D. (2004). *Quantum Entropy and its Use.* Springer Verlag, Berlin-Heidelberg.

[18] Petz, D. (1990). *An Invitation to the Algebra of Canonical Commutation Relations*, Leuven Notes in Mathematical and Theoretical Physics. Series A: Mathematical Physics Vol. 2 (Leuven University Press, Leuven).

[19] Petz, D. (2010). *, Quantum Information Theory and Quantum Statistics.* Springer.

[20] van der Vaart, A.W. (1998). *Asymptotic Statistics.* Cambridge University Press.

[21] Yamagata, K. (2011). Efficiency of quantum state tomography for qubits. *Int. J. Quant. Inform.,* **9** 1167.